

06/10/99

Docket No. 55293-B/JPW/ARC

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Respectable Assistant Commissioner for Patents  
Washington, D.C. 20231

June 10, 1999

1553 U.S. PTO  
09/329917  
06/10/99

S I R:

Transmitted herewith for filing are the specification and claims of the patent application of:

Carlos Cordon-Cardo  
Inventor(s) for

MARKERS FOR PROSTATE CANCER

Title of Invention

Also enclosed are:

☒ 13 sheet(s) of ☐ informal ☒ formal drawings.

☐ Oath or declaration of Applicant(s).

☐ A power of attorney

☐ An assignment of the invention to

☒ A Preliminary Amendment

☐ A verified statement to establish small entity status under 37 C.F.R. §1.9 and §1.27.

The filing fee is calculated as follows:

CLAIMS AS FILED, LESS ANY CLAIMS CANCELLED BY AMENDMENT

|   | NUMBER<br>FILED |   | NUMBER<br>EXTRA* |   | RATE            |                 | FEE |                 |                 |
|---|-----------------|---|------------------|---|-----------------|-----------------|-----|-----------------|-----------------|
|   |                 |   |                  |   | SMALL<br>ENTITY | OTHER<br>ENTITY |     | SMALL<br>ENTITY | OTHER<br>ENTITY |
| Total Claims  | 6   -20         | = | 0                | X | 9               | 18              | =   | \$ 0            | \$              |
| Independent<br>Claims   | 3   -3          | = | 0                | X | 39              | 78              | =   | \$ 0            | \$              |
| Multiple Dependent<br>Claims Presented: <u>      </u> Yes <u>  X  </u> No |                 |   |                  |   | 130             | 260             | =   | \$ 0            | \$              |
| *If the different in Col. 1 is<br>less than zero, enter "0" in<br>Col. 2  |                 |   |                  |   | BASIC FEE       |                 |     | 380             | 760             |
|   |                 |   |                  |   | TOTAL FEE       |                 |     | \$ 380          | \$              |

Applicant : Carlos Cordon-Cardo  
Serial No.: Not Yet Known  
Filed : Herewith

Letter of Transmittal  
Page 2

- ☒ A check in the amount of \$ 380.00 to cover the filing fee.
- ☐ Please charge Deposit Account No. \_\_\_\_\_ in the amount of \$ \_\_\_\_\_.
- ☒ The Commissioner is hereby authorized to charge any additional fees which may be required in connection with the following or credit any over-payment to Account No. 03-3125:
- ☒ Filing fees under 37 C.F.R. \$1.16.
- ☒ Patent application processing fees under 37 C.F.R. \$1.17.
- ☐ The issue fee set in 37 C.F.R. \$1.18 at or before mailing of the Notice of Allowance, pursuant to 37 C.F.R. \$1.311(b).
- ☒ Three copies of this sheet are enclosed.
- ☐ A certified copy of previously filed foreign application No. \_\_\_\_\_ filed in \_\_\_\_\_ on \_\_\_\_\_.
- ☐ Applicant(s) hereby claim priority based upon this aforementioned foreign application under 35 U.S.C. §119.
- ☒ Other (identify) Abstract 1 page: Claims 4 pages; One loose set of formal drawings; Express Mail Certificate of Mailing bearing Label No. EL278886800US, dated June 10, 1999.

Respectfully submitted,

Albert Wai Kit Chan

John P. White  
Registration No. 28,678  
Albert Wai-Kit Chan  
Registration No. 36,479  
Attorneys for Applicants  
Cooper & Dunham LLP  
1185 Avenue of the Americas  
New York, New York 10036  
(212) 278-0400

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant : Carlos Cordon-Cardo  
Serial No. : Not Yet Known  
(Continuation-In-Part of International  
Application No. PCT/US98/25483, filed  
December 1, 1998, which claims the  
benefit of U.S. Provisional Application  
No. 60/067,190, filed December 1, 1997)  
Filed : Herewith  
For : MARKERS FOR PROSTATE CANCER

1185 Avenue of the Americas  
New York, New York 10036  
June 10, 1999

Assistant Commissioner for Patents  
Washington, D.C. 20231

BOX: Patent Applications

SIR:

PRELIMINARY AMENDMENT

Please amend the above-identified application as follows:

In the claims:


Please cancel claims 4-5, and 9-18 without prejudice to applicant's right to pursue the subject matter in a future continuation or divisional application. Accordingly, claims 1-3, and 6-8 are pending in this application.

If a telephone interview would be of assistance in advancing prosecution of the subject application, applicant's undersigned attorney invites the Examiner to telephone at the number provided below.

Applicant : Carlos Cordon-Cardo  
Serial No.: Not Yet Known  
Filed : Herewith  
Page 2

No fee is deemed necessary in connection with the filing of this Preliminary Amendment. However, if any additional fee is required, authorization is hereby given to charge the amount of such fee to Deposit Account No. 03-3125.

Respectfully submitted,



John P. White  
Registration No. 28,678  
Albert Wai-Kit Chan  
Registration No. 36,479  
Attorneys for Applicant(s)  
Cooper & Dunham, LLP  
1185 Avenue of the Americas  
New York, New York 10036  
(212) 278-0400



*Application  
for  
United States Letters Patent*

**To all whom it may concern:**

Be it known that **Carlos Cordon-Cardo**

have invented certain new and useful improvements in

**MARKERS FOR PROSTATE CANCER**

of which the following is a full, clear and exact description.

MARKERS FOR PROSTATE CANCER

This application is a Continuation-In-Part application of International Application No. PCT/US98/25483, filed December 1, 1998, which claims the benefit of U.S. Provisional Application No. 60/067,190, filed December 1, 1997, the content of which are incorporated into this application by reference.

This invention was made in part with support under United States Government NIH Grant CA-DK-47650. Accordingly, the United States Government has certain rights in the invention.

Throughout this application, various references are referred to within parentheses. Disclosures of these publications in their entirety are hereby incorporated by reference into this application to more fully describe the state of the art to which this invention pertains. Full bibliographic citation for these references may be found at the end the specification, preceding the claims.

20

Background of the Invention

It has been postulated that the loss of function of a new family of negative cell cycle regulators, which act as cyclin-dependent kinase inhibitors and have been termed CKI, might lead to tumor development. CKIs fall into two families, Kip and Ink, on the basis of sequence homology. p27<sup>Kip1</sup> is implicated in G1 phase arrest by associating with multiple G1 cyclin-dependent kinases, abrogating their activity. However, no tumor-specific p27<sup>Kip1</sup> genomic mutations have been found in a large group of primary human cancers studied. More recently, it has been reported that proteasome-mediated degradation of p27 protein occurs during the cell cycle and that this degradation is

increased in a subset of breast and colon carcinomas of poor prognosis. **Purpose:** The present study was undertaken in order to assess for potential alterations of p27 expression in benign prostatic hyperplasia (BPH) and in a well characterized cohort of patients with prostatic cancer.

Inactivation of the p53 and RB tumor suppressor genes has been implicated in the development and progression of a number of different cancers (1,2). It has also been postulated that the loss of function of a new family of negative cell cycle regulators, which act as cyclin-dependent kinase inhibitors and have been termed CKI, might also lead to tumor development (3). CKIs fall into two families, Kip and Ink, on the basis of sequence homology (4). Kip family members include p21 (also known as WAF1, Cip1, or Sdi1) (5-7), p27<sup>Kip1</sup> (8-10) and p57<sup>Kip2</sup> (11,12). The Ink group includes four members: p16<sup>INK4A/MTS1/CDKN2</sup> (13), p15<sup>INK4B/MTS2</sup> (14), p18<sup>INK4C</sup> (15), and p19<sup>INK4D</sup> (16). p27 is a negative regulator implicated in G1 phase arrest by TGF $\beta$ , cell-cell contact, agents that elevate cyclic AMP, and the growth inhibitory drug rapamycin (17-21). p27 associates with multiple G1 cyclin-dependent kinases in non-proliferating cells, abrogating their activity (4, 8-10).

To assess its role as a potential tumor suppressor, the p27<sup>Kip1</sup> gene was mapped to 12p12-12p13.1 and no tumor-specific genomic mutations in a large group of primary human cancers were observed (22-24). More recently, it has been reported that proteasome-mediated degradation of p27 occurs during the cell cycle and that this degradation is increased in a subset of breast and colon carcinomas of poor prognosis (25-28). The present study was undertaken in order to assess for potential

alterations of p27 expression in BPH and in a well characterized cohort of patients with primary and metastatic prostatic cancer.

5 74 prostate carcinomas from primary and metastatic sites, representing different hormone sensitivities were analyzed. Normal prostatic tissues and cases of benign prostatic hyperplasia were also studied. In order to evaluate  
10 prostatic tissue of p27 null mice, eight 7 month old and six greater than 12 month old littermate pairs of wild-type and p27 knockout animals were used. Levels of expression and microanatomical localization of p27 protein and RNA transcripts were determined by immunohistochemistry and *in situ* hybridization with specific antibodies and probes,  
15 respectively. Comparative analyses between immunohistochemistry, immunoblotting and immunodepletion assays were also conducted in a subset of cases. Association between alterations in p27 expression and clinicopathological variables were evaluated using the  
20 two-tailed Fisher's exact test. Disease relapse-free survivals were evaluated using the Kaplan-Meier method and the Logrank test. Distinct anomalies in the expression of p27 in benign and malignant human prostate tissues are reported. The normal human prostate shows abundant amounts  
25 of p27 and high levels of p27 messenger in both epithelial and stroma cells. However, p27 protein and transcripts are almost undetectable in epithelial and stroma cells of BPH lesions. It is also reported that p27-null mice develop hypercellular prostatic glands which histologically  
30 resemble human BPH. Based on these findings we postulate that the loss of p27 expression in human prostate may be causally linked to BPH. Prostatic carcinomas can be categorized into two groups: those that contain detectable p27 protein and those that do not. In contrast to BPH,  
35 however, both groups of prostatic carcinomas contain

abundant p27 transcripts. Moreover, primary prostatic carcinomas displaying the p27-negative phenotype appear to be biologically more aggressive, based on their association with time to prostate specific antigen (PSA) failure following radical prostatectomy. These results support the postulate that BPH is not a premalignant lesion in the pathway of prostate cancer development. Data also suggest that prostatic carcinoma develops along two different pathways, one involving the loss of p27 and the other using other processes that circumvent the growth suppressive effects of p27.

Summary of the Invention

This invention provides a method for determining the aggressiveness of a prostate carcinoma comprising: (a) obtaining a sample of the prostate carcinoma; and (b) detecting the presence of p27 protein in the prostate carcinoma, the absence of p27 indicating that the prostate carcinoma is aggressive.

This invention also provides a method for diagnosing a benign prostate hyperplasia comprising: (a) obtaining an appropriate sample of the hyperplasia; and (b) detecting the presence of the p27 RNA, a decrease of the p27 RNA indicating that the hyperplasia is benign. In an embodiment, the above method further detects the protein expression of p27 wherein this additional step may be performed before or after the detection of the presence of the p27 RNA.

This invention provides a method for predicting the life-span of patient with prostate carcinoma comprising: (a) obtaining a sample of the prostate carcinoma; and (b) detecting the presence of p27 protein in the prostate carcinoma, the presence of the p27 protein indicating that the patient can live longer than the patient who are undetectable p27 protein.

This invention also provides a method for increasing the life-span of patient with prostate carcinoma comprising inducing the expression of p27 protein in the prostate carcinoma.

This invention provides a method for prolong life-span of patient with prostate carcinoma which comprises introducing a nucleic acid molecule having sequence encoding a p27 protein into the carcinoma cell under conditions permitting

expression of said gene so as to prolong the life-span of the patient with said prostate carcinoma. In an embodiment, the nucleic acid molecule comprises a vector. The vector includes, but is not limited to, an adenovirus  
5 vector, adeno-associated virus vector, Epstein-Barr virus vector, Herpes virus vector, attenuated HIV virus, retrovirus vector and vaccinia virus vector.

10 This invention provides a method for prolong life-span of patient with prostate carcinoma which comprises introducing an effective amount of p27 protein into the carcinoma cell so as to thereby prolong the life-span of the patient with said prostate carcinoma.

15 This invention provides a method for prolong life-span of patient with prostate carcinoma which comprises introducing an effective amount of a substance capable of stabilizing the p27 protein into the carcinoma cell so as to thereby prolong the life-span of the patient with said prostate  
20 carcinoma.

This invention provides a composition for prolong life-span of patient with prostate carcinoma which comprises an effective amount of a nucleic acid molecule having sequence  
25 encoding a p27 protein and a suitable carrier.

This invention provides a composition for prolong life-span of patient with prostate carcinoma which comprises an effective amount of the p27 protein and a suitable carrier.

30 This invention provides a composition for prolong life-span of patient with prostate carcinoma which comprises an effective amount a substance capable of stabilizing the p27 protein and a suitable carrier.

35

## Brief Description of the Figures

### Figures for the First Series of Experiments

5     **Figure 1.**

Histological analysis, immunohistochemistry, and *in situ* hybridization of human primary and metastatic prostatic carcinomas.

10     (A-C) Photomicrographs of primary prostatic carcinomas processed as follows: (A) Immunohistochemical staining against p27 of a prostatic intra-epithelial neoplastic (PIN) lesion; note the intense positive immunoreactivities observed in the nuclei of the tumor cells growing into the lumen. (B) Immunohistochemical staining against p27 of  
15     another PIN lesion showing dysplastic changes; note the intense positive immunostaining in the nuclei of normal epithelial cell and the low-to-undetectable staining of the tumor cells dissecting the gland and growing into the lumen. (C) Undetectable levels of p27 protein in an  
20     invasive primary prostatic carcinoma; note the staining of a normal gland trapped into the tumor.

(D-F) Photomicrographs of metastatic prostatic carcinomas processed as follows: (D) Immunohistochemical staining against p27 of a metastatic prostate carcinoma to lymph  
25     node; note the intense nuclear staining of both tumor cells and lymphocytes (cells in the germinal center display low p27 levels). (E) Immunohistochemical staining against p27 of another metastatic prostate carcinoma to lymph node; note the intense positive immunostaining in the nuclei of  
30     lymphocytes and the undetectable levels of p27 staining on the tumor cells. (F) Immunohistochemical staining against p27 of a metastatic prostate carcinoma to bone; note the positive immunoreactivities in the nuclei of osteoblasts and the lack of staining of tumor cells.

35     (F and G) Photomicrographs of a primary invasive prostatic



carcinoma processed as follows: (F) Low-to undetectable immunohistochemical staining against p27 in the tumor cells; note the staining of a normal gland trapped into the tumor. (G) *In situ* hybridization on a consecutive section from the case illustrated in panel (F) showing high mRNA levels of p27<sup>Kip1</sup> even in p27-negative tumor cells utilizing the anti-sense probe to p27<sup>Kip1</sup>. Original magnification (A) through (F) 400x.

10 Figure 2.

In certain prostatic carcinomas p27 protein is a functional cyclin-dependent kinase inhibitor. (A) Immunohistochemical staining correlates with the presence of p27 by immunoblotting. Tumors #1 and #2 were negative and tumor #3 positive for p27 protein expression, paralleling their IHC patterns. (B) Immunodepletion of p27 extracts. Extracts obtained from tumors #2 and #3 were subjected to sequential depletion with antibodies specific to p27 or a non-specific rabbit anti-mouse (RaM). Following depletion, the proteins in the supernatants were resolved and the presence of p27 determined by immunoblotting. (C) Depletion of p27 depletes heat stable cyclin-dependent kinase inhibitory activity. The supernatant shown in panel B was boiled and following clarification the soluble fraction was incubated with different amounts of recombinant cyclin E/CDK2 kinase and the degree of inhibition of cyclin E/CDK2 activity on histone H1 substrate was measured. The amount of each kinase used is shown in the panel and the bars are representative activities on an arbitrary scale. Depletion with either RaM or p27 specific antibodies did not affect the inhibitory activity of the p27 negative tumor; however, depletion of p27 from the positive tumor extract completely ablated the heat stable inhibitor activity.

**Figure 3.**

Recurrence-free proportion analysis of patients with primary prostate carcinoma (n=42) as assessed by time to detectable PSA. Patients who had PSA relapse were classified as failures, and patients with PSA relapse, or those who were still alive or died from other disease or lost to follow-up during the study period, were coded as censored. Time to relapse was defined as the time from date of surgery to the endpoint (relapse or censoring). Disease relapse-free survivals were evaluated using the Kaplan-Meier method and the Logrank test. A trend was observed between a p27 negative phenotype and early relapse (p=0.08).

**Figure 4.**

Histological analysis, immunohistochemistry, and *in situ* hybridization of human normal prostate and benign prostatic hyperplasia.

(A-C) Photomicrographs of consecutive sections of normal prostate tissue processed as follows: (A) Immunohistochemical staining against p27; intense positive immunoreactivities are observed in the nuclei of epithelial cells in the luminal side of the acinus, with decreased reactivities in the nuclei of basal and stroma cells. (B) *In situ* hybridization showing high mRNA levels of p27<sup>Kip1</sup> in both epithelial and stroma cells utilizing the anti-sense probe. (C) *In situ* hybridization utilizing the sense probe to p27<sup>Kip1</sup> showing lack of signals in both epithelial and stroma cells.

(D-F) Photomicrographs of consecutive tissue sections of a benign prostatic hyperplastic nodule processed as follows: (D) Immunohistochemical staining against p27; note the lack or almost undetectable levels of immunoreactivity observed in the nuclei of both epithelial and stroma cells in the luminal side of the acinus, with decreased

reactivities in the nuclei of basal and stroma cells. (E) *In situ* hybridization showing low-to-undetectable p27<sup>Kip1</sup> transcripts also in both epithelial and stroma cells utilizing the anti-sense probe; note the strong signal of the cellular inflammatory infiltrates that serve as an internal positive control. (F) *In situ* hybridization utilizing the sense probe to p27<sup>Kip1</sup> showing lack of signals in epithelial and stroma cells, as well as cellular inflammatory elements. Original magnifications: (A), (B) and (C) 1000x; (D), (E) and (F) 400x.

**Figure 5.**

Histopathological analysis of the prostatic tissues of 12 month old p27<sup>+/+</sup> (A) and p27<sup>-/-</sup> (B-D) mice. Photomicrographs of tissue sections of normal prostate samples processed as follows: (A) Hematoxylin and eosin staining of a prostate gland of a p27<sup>+/+</sup> mouse showing well defined acini of epithelial cells surrounded by a stroma containing few fibroblasts and poor in supportive connective tissue components. (B) Hematoxylin and eosin staining of a prostate gland of a p27<sup>-/-</sup> mouse showing multiple and complex glands and hypercellular acini of epithelial cells surrounded by fibromuscular stroma cells in a connective tissue displaying abundant supportive components. (C and D) Hematoxylin and eosin stainings of a prostate gland of a p27<sup>-/-</sup> mouse, high power details, illustrating the complexity of the glands and abundant fibromuscular stroma elements (C), as well as the hypercellularity of the acini (D). Original magnifications: (A) and (B) 200x; (C) and (D) 400x.

## Figures for the Second Series of Experiments

- Figure 6.** Photomicrographs of selected primary prostate carcinoma cases analyzed by immunohistochemistry utilizing mouse monoclonal antibodies PAB1801 (anti-p53, A), 2A10 (anti-mdm2, B), Ab-1 (anti-p21, C), and MIBI (anti-Ki67, D). A, p53 nuclear overexpression in tumor cells. Note the positive nuclear staining of tumor cells at a perineural invasion site (arrow). B, mdm2 nuclear overexpression in tumor cells. C, p21 nuclear overexpression in tumor cells. D, high Ki67 proliferative index. Note the intense Ki67 nuclear staining of tumor cells at a perineural invasion site (arrow).
- Figure 7.** Progression-free and survival curves for patients with primary prostate cancer. The Kaplan-Meier method was used to estimate overall disease free survival. The log-rank analysis was used to compare the different curves. A, progression was significantly reduced in patients with tumors displaying a p53-positive phenotype ( $P < 0.01$ ). B, progression was not related to mdm2 status. C, progression was significantly reduced in patients with tumors displaying a p21 positive phenotype ( $P = 0.0165$ ).
- Figure 8.** Diagrammatic representation of the p53-pathway (A), and alterations that may develop during tumor progression in prostate cancer (B). (A) p53 regulates the expression of several genes involved in cell cycle arrest (ie, p21) and apoptosis (ie, bax). p21 binds to heterodimeric protein kinases formed by cyclins and cyclin-dependent kinases (Cdk's), blocking phosphorylation of pRB/E2F1 complexes and abrogating S-phase entry. p53 also produces an autoregulatory feed back loop by transactivating mdm2. (B) Overexpression of mdm2 has been observed to occur in several tumor types, and it is

considered an oncogenic event. Upon binding to mdm2, p53 products are transcriptionally inactivated and triggered for degradation. This will release the G1 arrest imposed, in part, by p21 and abolish the apoptotic signals of the pathway. Thus, inactivation of p53 will favor proliferative activity, immortality, and development/accumulation of further DNA damage or mutations. The increased p21 expression observed in our study could be produced via growth factor signaling, which would also impact on cyclin D1 expression. The increment of p21 does not appear to be able to control the proliferative activity of tumor cells, as attested by the association of p21 positive phenotype and high Ki67 proliferative index. Taken together, mdm2 overexpression will inactivate the p53-pathway, while increased mitogenic activity will offset the RB-pathway. The mechanistic basis for this dual requirement stems, in part, from the deactivation of a p53-dependent cell suicide program that would normally be brought about as a response to unchecked cellular proliferation resulting from RB-deficiency.

#### Figures for the Third Series of Experiments

Figure 9A., Figure 9B., Figure 9C., Figure 9D., Figure 9E., and Figure 9F.

#### Figures for the Fourth Series of Experiments

Figure 10A-B. Immunohistochemistry and in situ hybridization of human benign prostatic hyperplasia (BPH). Consecutive sections of benign hyperplastic prostate tissue were processed as follows: A) Immunohistochemical staining of p16 is shown. Protein expression levels are undetectable in both epithelial and stromal components. B) In situ hybridization shows undetectable mRNA levels of p16

in both epithelial and stromal components when the antisense probe is used.

**Figure 11A-D.** Immunohistochemistry and in situ hybridization of human primary prostatic carcinomas. Consecutive sections of primary human prostate cancer tissue were processed as follows: A) Immunohistochemical staining of p16 is shown. Lack of immunoreaction noted in the nuclei and cytoplasm of both epithelial and stromal components. B) In situ hybridization reveals undetectable mRNA levels of p16 in both epithelial and stromal components when the antisense probe is used. (C-D) Histologic analysis, immunohistochemistry, and in situ hybridization of human primary prostatic carcinoma showing p16 overexpression. Consecutive sections of primary human prostate cancer tissue were processed as follows: C) Immunohistochemical staining of p16 is shown. Note strong brown immunoreaction observed in the nuclei of cells. Faint cytoplasmic staining is noted as well. D) In situ hybridization shows high mRNA levels of p16 in epithelial cells when the antisense probe is used. A normal gland (see pointer) serves as an internal negative control in both the immunohistochemical analysis in Figure 11C and also the in situ hybridization analysis in Figure 11D.

**Figure 12.** Kaplan-Meier curves, using the log rank test, stratified by p16 groups (group A or group B) of patients with primary prostate carcinoma (n=88) as assessed by time to detectable PSA level post prostatectomy. Time to relapse was defined as the time from the date of surgery to the time of PSA elevation after surgery. The median time to relapse for group A has not been reached. The median time to relapse for group B was 46.25 months. Patients who had PSA relapse were classified as having treatment

4. 3

Detailed Description of the Invention

5 This invention provides a method for determining the aggressiveness of a prostate carcinoma comprising: (a) obtaining a sample of the prostate carcinoma; and (b) detecting the presence of p27 protein in the prostate carcinoma, the absence of p27 indicating that the prostate carcinoma is aggressive.

10 This invention also provides a method for diagnosing a beign prostate hyperplasia comprising: (a) obtaining an appropriate sample of the hyperplasia; and (b) detecting the presence of the p27 RNA, a decrease of the p27 RNA indicating that the hyperplasia is beign. In an  
15 embodiment, the above method further detects the protein expression of p27 wherein this additional step may be performed before or after the detection of the presence of the p27 RNA.

20 This invention provides a method for predicting the life-span of patient with prostate carcinoma comprising: (a) obtaining a sample of the prostate carcinoma; and (b) detecting the presence of p27 protein in the prostate carcinoma, the presence of the p27 protein indicating that  
25 the patient can live longer than the patient who are undetectable p27 protein.

This invention also provides a method for increasing the life-span of patient with prostate carcinoma comprising  
30 inducing the expression of p27 protein in the prostate carcinoma.

This invention provides a method for prolong life-span of patient with prostate carcinoma which comprises introducing  
35 a nucleic acid molecule having sequence encoding a p27



protein into the carcinoma cell under conditions permitting expression of said gene so as to prolong the life-span of the patient with said prostate carcinoma. In an embodiment, the nucleic acid molecule comprises a vector.

5 The vector includes, but is not limited to, an adenovirus vector, adeno-associated virus vector, Epstein-Barr virus vector, Herpes virus vector, attenuated HIV virus, retrovirus vector and vaccinia virus vector.

10 Methods to introduce a nucleic acid molecule into cells have been well known in the art. Naked nucleic acid molecule may be introduced into the cell by direct transformation. Alternatively, the nucleic acid molecule may be embedded in liposomes. Accordingly, this invention  
15 provides the above methods wherein the nucleic acid is introduced into the cells by naked DNA technology, adenovirus vector, adeno-associated virus vector, Epstein-Barr virus vector, Herpes virus vector, attenuated HIV vector, retroviral vectors, vaccinia virus vector,  
20 liposomes, antibody-coated liposomes, mechanical or electrical means. The above recited methods are merely served as examples for feasible means of introduction of the nucleic acid into cells. Other methods known may be also be used in this invention.

25 This invention provides a method for prolong life-span of patient with prostate carcinoma which comprises introducing an effective amount of p27 protein into the carcinoma cell so as to thereby prolong the life-span of the patient with  
30 said prostate carcinoma.

This invention provides a method for prolong life-span of patient with prostate carcinoma which comprises introducing an effective amount of a substance capable of stabilizing  
35 the p27 protein into the carcinoma cell so as to thereby

prolong the life-span of the patient with said prostate carcinoma. Such substance may be either inhibiting the protease which degrade the p27 protein or it may interact with p27 in such a way that the protein will be resistant to degradation. By administering such substance into the cell, the effective amount of p27 protein will be increased.

This invention provides a composition for prolong life-span of patient with prostate carcinoma which comprises an effective amount of a nucleic acid molecule having sequence encoding a p27 protein and a suitable carrier.

As used herein, the term "suitable carrier" encompasses any of the standard carriers. The composition may be constituted into any form suitable for the mode of administration selected. Compositions suitable for oral administration include solid forms, such as pills, capsules, granules, tablets, and powders, and liquid forms, such as solutions, syrups, elixirs, and suspensions. Forms useful for parenteral administration include sterile solutions, emulsions, and suspensions.

This invention provides a composition for prolong life-span of patient with prostate carcinoma which comprises an effective amount of the p27 protein and a suitable carrier.

This invention provides a composition for prolong life-span of patient with prostate carcinoma which comprises an effective amount a substance capable of stabilizing the p27 protein and a suitable carrier.

This invention provides a method for determining the rate of proliferation of a prostate cancer comprising: (a) obtaining a sample of the prostate cancer; and (b)

detecting the presence of p21 protein in the prostate cancer, the presence of p21 indicating that the prostate cancer will have a high proliferation rate.

5 This invention also provides a method for determining the rate of proliferation of a prostate cancer comprising: (a) obtaining a sample of the prostate cancer; and (b) detecting the mdm2 expression in the prostate cancer, the overexpression of mdm2 indicating that the  
10 prostate cancer will have high proliferation rate.

This invention provides a method for determining whether a prostate cancer would be metastatic comprising: (a) obtaining a sample of the prostate cancer; and (b) detecting  
15 the level of cyclin D1 expression in the prostate cancer, the overexpression of cyclin D1 indicating that the prostate cancer will be metastatic. In an embodiment, the prostate cancer is metastatic to bone.

20 This invention provides a method for determining the tumor recurrence in prostate cancer comprising: (a) obtaining a sample of the prostate cancer; and (b) detecting the expression of the cyclin-dependent kinase inhibitor p16 in the prostate cancer, the overexpression of p16 indicating  
25 that the prostate cancer will have high tumor recurrence.

This invention will be better understood from the Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods  
30 and results discussed are merely illustrative of the invention as described more fully in the claims which follow thereafter.

## EXPERIMENTAL DETAILS

### Experimental Details for First Series of Experiments

#### 5 MATERIALS AND METHODS

Patient Characteristics and Tissues. A cohort of 74 patients with prostatic carcinoma were evaluated. Tissues were obtained from the Department of Pathology, Memorial Sloan-Kettering Cancer Center, New York. Samples were formalin-fixed, paraffin-embedded tissue specimens. Forty-two primary prostate adenocarcinoma specimens were evaluated, as well as 9 metastases to lymph node and 23 metastases to bone. Normal prostatic tissue and/or areas of benign prostatic hyperplasia adjacent to tumor were observed in the majority of the primary cases studied. These tissues were also analyzed as part of the study. In addition, 10 pairs of frozen normal and tumor prostate tissues were utilized for antibody titration, as well as comparative analyses between immunohistochemistry, immunoblotting and immunodepletion assays (see below). Representative hematoxylin-eosin stained sections were examined to evaluate the histopathological characteristics of the lesions to be analyzed, including the ratio of normal-to-tumor content for microdissection techniques.

25 In order to evaluate prostatic tissue of p27 null mice, eight 7 month old and six greater than 12 month old littermate pairs of wild-type and p27 knockout animals were used. Tissues were dissected, weighted and processed for histology by formalin fixation and paraffin embedding. Tissue sections were cutted and stained with hematoxylin-eosin for histologic analysis. All sections were utilized to count the number of acini per gland, a process that was conducted utilizing magnifications of 35 200x.

**Antibodies and Immunohistochemistry.** The following well characterized antibodies and corresponding final working dilutions were used for the present study: monoclonal antibody p27/Kip1 (Ab-2, Oncogene Science, Boston, MA - 0.1 ug/ml final concentration) and anti-p27 affinity purified rabbit antiserum (1:500 dilution). A non-immune rabbit serum and mouse monoclonal antibody M1gS-Kp1 were used as negative controls at similar working dilutions.

Deparaffinized sections were treated with 3% H<sub>2</sub>O<sub>2</sub> in order to block endogenous peroxidase activity. Sections were subsequently immersed in boiling 0.01% citric acid (pH 6.0) in a microwave oven for 15 minutes to enhance antigen retrieval, allowed to cool, and incubated with 10% normal horse or normal goat sera to block non-specific tissue immunoreactivities. Primary antibodies were then incubated overnight at 4°C. Biotinylated horse anti-mouse IgG antibodies (Vector Laboratories, Burlingame, CA - 1:500 dilution) or goat anti-rabbit antibodies (Vector Laboratories - 1:800 dilution) were applied for 1 hour, followed by avidin-biotin peroxidase complexes for 30 minutes (Vector Laboratories - 1:25 dilution). Diaminobenzidine was used as the final chromogen and hematoxylin was used as the nuclear counterstain. Nuclear immunoreactivities were classified as a continuum data (undetectable levels or 0% to homogeneous staining or 100%). Tumors were grouped into two categories defined as follows: negative (0% or undetectable staining to <20% nuclear immunoreactivity in tumor cells), and positive (neoplasms with ≥20% tumor cells with nuclear staining) (see statistical section).

**Probes and *In Situ* Hybridization.** Digoxigenin-labeled probes were used for in situ hybridization and 1 ug of recombinant plasmid pCR<sup>TM</sup>II (Invitrogen, San Diego, CA),

containing the full length human p27 gene (gift of Dr. M. Pagano, New York University School of Medicine, NY) was linearized by BamHI and XbaI to generate antisense and sense transcripts. Riboprobes were generated with T7 and SP6 polymerase for 2 hours at 37°C in 1X transcription buffer (Boehringer Mannheim, Indianapolis, IN), 20 U of RNase inhibitor, 1 mmol/L each of ATP, GTP, CTP, 6.5 mmol/L UTP and 3.35 mmol/L digoxigenin-UTP. Deparaffinized tissue sections were rinsed in water and PBS for 10 minutes. The slides were digested with Proteinase K (50ug/ml) for 18 minutes at 37°C in PBS, and post-fixed at 4 °C in a freshly prepared solution of 4% paraformaldehyde in PBS for 5 minutes. Prehybridization was done for 30 minutes at 45 °C in 50% formamide and 2XSSC. The hybridization buffer consisted of 50% deionized formamide (v/v), 10% dextran sulphate (50% stock solution), 2XSSC (20X stock solution), 1% SDS (10% stock solution), and 0.25 mg/ml of herring sperm DNA (10 mg/ml). Hybridization was performed overnight at 45 °C applying 10 pmol/L digoxigenin-labeled riboprobe in 50 ul of hybridization buffer per section under a coverslip. The coverslips were removed and the slides were washed in pre-warmed 2XSSC for 20 minutes at 60 °C twice, followed by washes in pre-warmed 0.5XSSC and 0.01XSSC at 60 °C for 20 minutes, respectively. After these washes the slides were incubated in normal sheep serum diluted in buffer pH 7.5 and successively in the same buffer with antibody anti-digoxigenin-AP (Boehringer Mannheim, Indianapolis, IN) at dilution of 1:1500 for 1 hour at room temperature. The visualization was accomplished by nitro-blue tetrazolium 5-bromo-4-chloro-3-indoylphosphate. The slides were counterstained with methyl green and mounted.

**Immunoblotting and Immunodepletion Assays.** Proteins were extracted from three OCT-embedded prostatic carcinomas and

resolved on polyacrylamide gels for immunoblotting with p27-specific antibodies. Extracts obtained from p27 positive and negative tumors were subjected to sequential depletion with antibodies specific to p27 or a non-specific rabbit anti-mouse (RaM). Following depletion, the proteins in the supernatants were resolved and the presence of p27 determined by immunoblotting. Aliquots of these supernatants were briefly boiled and following clarification the soluble fraction was incubated with different amounts of recombinant cyclin E/CDK2 kinase and the degree of inhibition of cyclin E/CDK2 activity on histone H1 substrate was measured.

**Statistical Methods.** The statistical analyses were conducted as follows. For alterations of the p27, we divided patients into two groups: p27 negative (0% or no immunohistochemical staining to <20% tumor cells displaying nuclear reactivities) or p27 positive ( $\geq 20\%$  tumor cells with nuclear immunostaining with IHC). The data analyses were conducted to explore the relationship between p27 alterations and clinicopathological variables such as presentation (primary, lymph node metastases, and bone metastases), clinical stage (B, C, D), total Gleason score (6 or less versus 7 or more), and hormonal status (naïve versus androgen-independent) in a total 74 patients. For 42 patients with primary prostate cancer who underwent radical prostatectomy, further analysis was conducted to evaluate the relationship between p27 alterations and clinical variables, including those described above and PSA relapse (yes and no). Two-tail Fisher's exact test was utilized to assess these associations and two tailed p-values were employed as a significant level (29). The FREQ procedure in SAS was used in this study (30). In the analysis of disease relapse-free survival, patients who had PSA relapse were classified as lost failures, and patients

5



## EXPERIMENTAL RESULTS AND DISCUSSION

### 5      **Experimental Results and Discussion for the First Series of Experiments**

To determine whether loss of p27 expression was a common feature in prostate cancer, we analyzed 74 prostate carcinomas from primary and metastatic sites, representing  
10      different hormone sensitivities. Included were 42 hormone-naïve primary tumors, some with associated prostatic intraepithelial neoplastic (PIN) lesions, and 32 metastatic carcinomas from lymph node tumors (n=9) and bone metastases (n=23). Thirteen of these metastatic  
15      lesions were from hormone-naïve cases, while the remaining 19 metastases were obtained after hormonal treatment. PIN lesions displaying a cribriform or pseudopapillary pattern expressed high levels of p27 protein (Figure 1A) and were associated with p27-positive invasive prostatic carcinomas.  
20      In contrast, PIN lesions displaying a flat growth pattern had low to undetectable p27 levels (Figure 1B) and were associated with p27-negative invasive tumors. Of the invasive primary prostatic carcinomas studied, 12 of 42 (28.5%) cases had an intense nuclear immunoreactive p27  
25      pattern in the malignant cells (data not shown). The remaining 30 (71.5%) primary neoplasms displayed altered patterns of expression: 12 cases had undetectable p27 levels (Figure 1C), while 18 cases had a heterogeneous pattern of expression (data not shown). In metastatic  
30      lesions, 7 of 32 (21.9%) showed intense p27 nuclear immunostaining in most tumor cells (Figure 1D). The remaining 25 (78.1%) metastatic lesions had either heterogeneous (data not shown) or undetectable nuclear expression of p27 (Figures 1E and 1F). Interestingly, all  
35      but one of the nine patients with hormone-independent bone

lesions displayed altered p27 expression. Four of these 9 cases had undetectable p27 protein expression (Figure 1F), 4 cases had heterogeneous patterns of p27 expression ranging from 30% to 40% tumor cells with weak positive staining, and one case displayed 80% positive tumor cells. However, high levels of p27<sup>Kip1</sup> mRNA, as determined by in situ hybridization to a p27 cDNA probe, were found in all tumors even when the lesions displayed undetectable levels of p27 protein (Figures 1G and 1H).

In the group of tumors that expressed p27, we next determined if the p27 protein was inactivated. To accomplish this we extracted protein from fresh frozen samples and measured the heat stable Cdk inhibitory activity, using cyclin E/CDK2 as a substrate, remaining in extracts following depletion with p27-specific antibodies as described previously (17) (Figure 2). Depletion of p27 protein was confirmed by immunoblotting. As expected, the depletion of extracts derived from p27 negative tumors did not affect the heat stable inhibitory activity, nor did depletion of p27 positive tumor extract with a non-specific rabbit-anti-mouse immunoglobulin. However, depletion of extracts derived from p27 positive tumors with the p27-specific antibody completely removed the inhibitory activity, indicating that p27 was functional as a Cdk inhibitor in these samples.

Taken together, these data suggest that prostatic carcinomas develop along two different pathways, one involving the loss of p27 and the other using alternative processes that may circumvent the growth suppressive effects of p27. In order to determine if these distinct pathways of prostate tumorigenesis correlate with clinical parameters, as reported for other tumor types (25-28), associations between p27 immunostaining, stage, total

Gleason score, and hormonal status of the tumor were assessed. No associations between detectable versus undetectable p27 protein, Gleason score (6 or less versus 7 or more), or hormonal status (naïve versus androgen-independent) were observed. To assess disease aggressiveness, we evaluated the time to PSA failure, the most sensitive indicator of success or failure following radical prostatectomy, in patients treated for localized disease. Only patients who had an undetectable PSA level after surgery, an indication that the resection was complete, were considered. A trend toward an association was observed between a p27 negative phenotype and early relapse ( $p=0.08$ ) (Figure 3). This difference did not reach statistical significance due to the limited sample size of the cohort analyzed. Supporting this concept is the fact that in a multivariate proportional hazards analysis, after controlling for stage and Gleason score, p27 status still was the strongest factor in predicting PSA relapse ( $p=0.07$ ).

These data suggest extending the characterization of p27 expression to normal prostate and benign prostatic hyperplasia. In the normal human prostate, abundant amounts of p27 protein were detected in the ductal and acinar cells, mainly luminal elements, as well as stroma cells using immunohistochemistry. Epithelial cells displayed a strong nuclear immunostaining signal (Figure 4A). Likewise, both epithelial and stroma cells expressed abundant p27 transcripts (Figures 4B and 4C), as detected by in situ hybridization. Strikingly, in 12 cases of BPH p27 expression was low to undetectable in epithelial and stroma cells in the hyperplastic nodules. Immunohistochemical staining revealed low to undetectable immunoreactivities in both epithelial and fibromuscular cells in the hyperplastic nodules (Figure 4D). This

contrasts with the strong p27 nuclear immunostaining phenotype observed in the normal prostate. Likewise, p27 mRNA transcript levels were low to undetectable on consecutive sections of BPH by in situ hybridization (Figure 4E and 4F). In some of these BPH tissue samples we found areas of basal cell hyperplasia. These cellular elements also had low to undetectable amounts of p27 protein and transcripts (data not shown). Nevertheless, in the non-hyperplastic regions of these same BPH samples, normal ductal and acinar epithelial cells, as well as stroma elements, showed high levels of p27 expression. These results indicate that in the development of BPH, p27 transcription may be down-regulated. This finding was quite unexpected as this gene product is generally regulated at post-transcriptional levels (35-37), although members of the nuclear hormone receptor superfamily are suggested to regulate p27<sup>Kip1</sup> mRNA levels (38).

The targeted deletion of the p27 locus in a murine model was recently reported (39-41). p27 deficient mice are viable and display organomegaly, increased body size and female infertility. These anomalies could not be attributed to a defect of the growth hormone/IGF-1 axis, rather, they resulted from excess proliferation prior to withdrawal of cells into a terminally differentiated state (39). No increased incidence of spontaneous tumors was observed; however, many p27-null mice developed a pituitary hyperplasia reminiscent of adenoma in the intermediate lobe. These data suggest that p27 deficiency leads to hyperplasia in many tissues and organs. The high frequency of benign prostatic hyperplasia (BPH) in men and the alterations on p27 expression in that condition suggested a parallel to p27 deficiency. Previous reports of histopathological analyses of p27 null mice did not include the prostate (39-41). We next set up to determine

the morphologic characteristics of the prostate gland in p27 deficient animals. Comparing the total mean prostate weights of 7 month old age-matched p27+/+ (n=8) and p27-/- (n=8) mice, the differences were not significant [mean +/- SD: 80.6 mg (+/- 8.6 mg) and 90.1 mg (+/- 13.3 mg), respectively (p=0.1)]. However, the mean acini counts of the total gland in these groups were significantly different [mean +/- SD: 50.4 (+/- 8.5) and 74.9 (+/- 8.9), respectively (p<0.01)]. A similar relationship was observed in the mean total prostate weights of the old (greater than 12 months) p27+/+ (n=6) and p27-/- (n=6) mice [mean +/- SD: 114.0 mg (+/- 18.5 mg) and 119.0 mg (+/- 26.8 mg), respectively (p=0.7)], and the mean acini counts [mean +/- SD: 54.7 (+/- 6.5) and 73.8 (+/- 5.3), respectively (p<0.01)]. The significant increase in the number of acini in both young and old p27 deficient mice was associated with histopathological differences that became more accentuated in the elderly group. The hyperplastic prostate of the older p27-/- mice showed enlarged glands, development of hypercellular acini of epithelial cells, and an increase in fibromuscular stroma cells (Figure 5). These histological changes are reminiscent of BPH in humans and support the hypothesis that the loss of p27 expression in human prostate may be causally linked to BPH.

It has been suggested that BPH and malignant prostate growth share a common pathway because they commonly coexist and demonstrate androgen dependency (42-44). However, this relationship remains unclear since BPH tends to develop in the transition zone, while the majority of carcinomas develop in the peripheral zone (45-48). Results from the present study reveal that, unlike in the BPH lesions, prostatic carcinoma cells regulate p27 expression at the post-transcriptional level. Taken together these data

support the postulate that BPH is not a premalignant lesion in prostate cancer development.

Coordinate inactivation of the pathways involving the p53 and RB genes appears to be an essential requirement for the genesis of most human cancers. However, both p53 mutations and RB alterations are reported to be late and uncommon events in prostate tumor progression (49-52). Contrary to these results, data from this study indicate that inactivation of p27 is a frequent and early event in some prostate cancers. It is thus our working hypothesis that p27 represents another pathway of tumor suppression in certain human tumors, prostate cancer being a paradigm in which this concept could be further tested.

In summary, data from this study suggest that p27<sup>Kip1</sup> gene ablation in the mouse causes a pronounced prostatic hyperplasia, and that the loss of p27 expression in human prostate may be causally linked to BPH. In addition, data from this study suggest that prostatic carcinoma develops along two different pathways, one involving the loss of p27 and the other using alternative processes that circumvent the growth suppressive effects of p27. These phenotypes can be identified as early as in the PIN stage. Moreover, primary prostatic carcinomas displaying the p27-negative phenotype appear to be biologically more aggressive, based on their association with time to PSA failure following radical prostatectomy while controlling for other variables. The consistent alteration of p27 expression observed in all androgen-independent metastatic lesions suggests an association with tumor progression, which may be the result of the metastatic process itself. Alternatively, it may be postulated that p27 positive tumors are more sensitive to androgen ablation, the primary treatment of metastatic disease. Finally, two dissimilar

mechanisms appear to be involved in the loss of p27 expression in BPH versus a subset of prostatic carcinomas. p27<sup>Kip1</sup> mRNA levels are extensively reduced in BPH, whereas p27 proteins are diminished to undetectable levels in some

5 prostatic carcinomas despite detectable p27 mRNA as the result of a post-transcriptional event. These results support the postulate that BPH is not a premalignant lesion in the pathway of prostate cancer development.

REFERENCES FOR THE FIRST SERIES OF EXPERIMENTS

1. Hartwell, L.H. & Kastan, M.B. (1994) *Science* **266**, 1821-1828.
- 5 2. Cordon-Cardo, C. (1995) *Am. J. Path.* **147**, 545-560.
3. Sherr, C.J. (1996) *Science* **274**, 1672-1677.
4. Sherr, C.J. & and Roberts, J.M. (1995) *Genes Dev* **9**,  
10 1149-1163.
5. Harper, J.W., Adami, G.R., Wei, N., Keyomarsi, K. & Elledge, S.J. (1993) *Cell* **75**, 805-816.
- 15 6. El-Deiry, W.S., Tokino, T., Velculescu, V.E., Levy, D.B., Parsons, R., Trent, J.M., Lin, D., Mercer, W.E., Kinzler, K.W. & Vogelstein, B. (1993) *Cell* **75**, 817-825.
- 20 7. Xiong, Y., Hannon, G.J., Zhang, H. Casso, D., Kobayashi, R. & Beach, D. (1993) *Nature* **366**, 701-704.
8. Polyak, K., Kato, J-Y., Solomon, M.J., Sherr, C.J., Mssague, J., Roberts, J.M. & Koff, A. (1994) *Genes Dev*  
25 **8**, 9-22.
9. Polyak, K., Lee, M-H., Erdjument-Bromage, H., Koff, A., Roberts, J.M., Tempst, P. & Massague, J. (1994) *Cell* **78**, 59-66.
- 30 10. Toyoshima, H. & Hunter, T. (1994) *Cell* **78**, 67-74.
11. Lee, M-H., Reynisdóttir, I. and Massague, J. (1995) *Genes Dev.* **9**, 639-649.  
35



12. Matsuoka, M., Edwards, M.C., Bai,, C., Parker, S., Zhang, P., Baldini, A., Harper, W. & Elledge, S.J. (1995) *Genes Dev.*, **9**, 650-662.
- 5 13. Serrano, M., Hannon, G.J. & Beach, D. (1993) *Nature* **366**, 704-707.
14. Hannon, G.J. & Beach, D. (1994) *Nature* **371**, 257-261.
- 10 15. Guan, K-L., Jenkins, C.W., Li, Y., Nichols, M.A., Wu, X., O'Keefe, C.L., Matera, A.G. & Xiong, Y. (1994) *Genes Dev* **8**, 2939-2952.
- 15 16. Chan, F.K.M., Zhang, J., Cheng, L., Shapiro, D.N. & Winoto, A. (1995) *Mol. Cell Biol.* **15**, 2682-2688.
17. Polyak, K., Kato, J-Y., Solomon, M.J., Sherr, C.J., Massague, J., Roberts, J.M. & Koff, A. (1994) *Genes Dev* **8**, 9-22.
- 20 18. Kato, J.Y., Matsuoka, M., Polyak, K., Massague, J. & Sherr, C.J. (1994) *Cell* **79**, 487-496.
19. Coats, S., Flanagan, W.M., Nourse, J. & Roberts, J.M. (1996) *Science* **272**, 877-880.
- 25 20. Luo, Y., Marx, S.O., Kiyokawa, H., Koff, A., Massague, J. & Marks, A.R. (1996) *Mol Cell Biol* **16**, 6744-6751.
- 30 21. Millard, S.S., Yan, J., Nguyen, J., Pagano, M., Kiyokawa, H. & Koff, A. (1997) *J Biol Chem* **272**, 7093-7098.
22. Ponce-Castaneda, M.V., Lee, M-H., Latres, E., Polyak, K., Lacombe, L., Montgomery, K., Mathew, S., Krauter,
- 35

K., Sheinfeld, J., Massague, J. & Cordon-Cardo, C.  
(1995) *Cancer Res* 55, 1211-1214.

- 5        23. Pietenpol, J.A., Bohlander, S.K., Sato, Y., Rowley,  
J.D., Papadopoulos, N., Liu, B., Friedman, C., Trask,  
B.J., Roberts, J.M., Kinzler, K.W., Vogelstein, B.  
(1995) *Cancer Res* 55, 1206-1210.
- 10       24. Bullrich, F., MacLachlan, T.K., Sang, N., Druck, T.,  
Veronese, M.L., Allen, S.L., Chiorazzi, N., Koff A.,  
Huebner, K., Croce, C.M. & Giordano, A. (1995) *Cancer  
Res* 55, 1199-1205.
- 15       25. Loda, M., Cukor, B., Tam, S.W., Lavin, P., Fiorentino,  
M., Draetta, G.F., Jessup, J.M., & Pagano, M. (1997)  
*Nature Med.* 3, 231-234.
- 20       26. Porter, P.L., Malone, K.E., Heagerty, P.J., Alexander,  
G.M., Gatti, L.A., Firpo, E.J., Daling, J.R. &  
Roberts, J.M. (1997) *Nature Med.* 3, 222-225.
- 25       27. Catzavelos, C., Bhattacharya, N., Ung, Y.C., Wilson,  
J.A., Roncari, L., Sandhu, C., Shaw, P., Yeger, H.,  
Morava-Protzner, I., Kapusta, L., Frannssen, E.,  
Pritchard, K.I. & Slingerland, J.M. (1997) *Nature  
Med.* 3, 227-230.
- 30       28. Tan, P., Cady, B., Wanner, M., Worland, P., Cukor, B.,  
Magi-Galluzzi, C., Lavin, P., Draetta, G., Pagano, M.  
& Loda, M. (1997) *Cancer Res.* 57, 1259-1263.
29. Mehta C.R. & Patel, N.R. (1983) *J Am Stat Assoc* 78,  
427-434.

30. SAS Institute Inc, SAS/STAT User Guide, version 6 (1990).
- 5 31. Kaplan, E.L & Meier, P. (1958) *J Am Stat Assoc* 53, 457-481.
32. Peto, R., Pike, M.C., Armitage, P., Breslow, N.E., Cox, D.R., Howard, S.V., Mantel, N., McPherson, K., Peto, J. & Smith, P.G. (1977) *Br J Cancer* 35, 1-39.
- 10 33. Cox, D.R., (1972) *J R Statist Soc* 34, 187-220.
34. Cox, D.R. (1975) *Biometrika* 62, 269-279.
- 15 35. Pagano, M., Tam, S.W., Theodoras, A.M., Beer-Romero, P., Del Sal, G., Chau, V., Yew, P.R., Draetta, G.F. & Rolfe, M. (1995) *Science* 269, 682-685.
36. King, R.W., Deshaies, R.J., Peters, J-M. & Kirschner, M., (1996) *Science* 274, 1652-1659.
- 20 37. Hengst, L. & Sherr, S.I. (1996) *Science* 271, 1861-1864.
38. Liu, M., Lee, M.H., Cohen, M., Bommakanti, M. & Freedman, L.P. (1996) *Genes & Dev* 10, 142-153.
- 25 39. Kiyokawa, H., Kineman, R.D., Manova-Todorova, K.O, Soares, V.C., Hoffman, E.S., Ono, M., Khanam, D., Hayday, A.C., Frohman, L.A. & Koff, A. (1996) *Cell* 85, 721-732.
- 30 40. Nakayama, K., Ishida, N., Shirane, M., Inomata, A., Inoue, T., Shishido, N., Horii, I., Loh, D.Y. & Nakayama, K-I. (1996) *Cell* 85, 707-720.
- 35

41. Fero, M.L., Rivkin, M., Tasch, M., Porter, P., Carow, C.E., Firpo, E., Polyak, K., Tsai, L-H., Broudy, V., Perlmutter, R.M., Kaushansky, K. & Roberts, J.M. (1996) *Cell* 85, 733-744.
- 5
42. Scher, H., Steinieck, G. & Kelly, W.K. (1995) *Urology* 46, 142-148.
43. Oesterling, J.E. (1996) *Prostate* (Suppl) 6, 67-73.
- 10
44. Linehan, W., Cordon-Cardo, C., Isaacs, W., Eds. DeVita, V.T., Hellman, S, Rosenberg, S.A. (J.B. Lippincott Company, Philadelphia, 1997).
- 15
45. Bostwick, D.G. (1996) *Cancer* 78, 330-336.
46. Parkinson, M.C. (1995) *Histopathol* 27, 301-307.
47. Bostwick, D., Pacelli, A., Lopez-Beltran, A. (1996) *Prostate* 29, 117-134.
- 20
48. Lipski, B.A., Garcia, R.L. & Brawer, M.K. (1996) *Semin Urol Oncol* 14, 149-155.
- 25
49. Aprikian, A.G., Sarkis, A.S., Fair, W.R., Zhang, Z-F., Fuks, Z. & Cordon-Cardo, C. (1994) *J Urol* 151, 1276-1280.
50. Brooks, J.D., Bova, G.S., Marshall, F.F. & Isaacs, W.B. (1993) *J Urol* 150, 1278-1283.
- 30
51. Dinjens, W.N., van der Weiden, M.M., Schroeder, F.H., Bosman,, F.T. & Trapman, J. (1994) *Int J Cancer* 56, 630-633.
- 35

52. Grignon, D.J., Caplan, R., Sarkar, F.H., Lawton, C.A., Hammond, E.H., Pilepich MV, Forman JD, Mesic J, Fu KK, Abrams RA, Pajak TF, Shipley WU & Cox JD. (1997) *J Natl Cancer Inst* 89, 158-165.

SECOND SERIES OF EXPERIMENTS

To determine the potential role of p53 inactivation in prostate cancer, we studied a well characterized cohort of 86 patients treated with radical prostatectomy. We analyzed patterns of p53, mdm2, and p21/WAF1 expression by immunohistochemistry. Results were then correlated with clinicopathological parameters of poor outcome, including time to PSA relapse. In addition, data were also correlated with proliferative index, as assessed by Ki67 antigen detection. p53 positive phenotype, defined as identification of nuclear immunoreactivity in >20% tumor cells, was observed in 6 of 86 cases (7%). An association was observed between p53 positive phenotype and decreased time to PSA relapse ( $P < 0.01$ ). mdm2 positive phenotype, defined as  $\geq 20\%$  tumor cells displaying nuclear immunoreactivity, was observed in 28 of 86 cases (32.5%). mdm2 positive phenotype was found to be associated with advanced stage ( $P = 0.009$ ). p21 positive phenotype, defined as >5% tumor cells with nuclear immunoreactivity, was observed in 28 of 86 cases (32.5%). An association was observed between p21 positive phenotype and high Ki67 proliferative index ( $P = 0.002$ ). Patients with p21 positive phenotype had a significant association with decreased time to PSA relapse ( $P = 0.0165$ ). In addition, a significant association was found between p21 positive phenotype and co-expression of mdm2 ( $P < 0.01$ ). Forty-three of 86 cases (50%) were found to have one or more alterations, and patients with any alteration were found to have a higher rate of PSA relapse ( $P < 0.01$ ). It is our hypothesis that a pathway of prostate cancer progression involves p53 inactivation caused by mdm2 overexpression, and that p21 transactivation in this setting is due to an alternative signaling system rather than through a p53-dependent mechanism.

p53 responds to different forms of cellular stress by targeting and activating genes involved in growth arrest and cell death. A target of p53-induced transcription is the p21/WAF1 gene, which encodes a cyclin-dependent kinase inhibitor (1). In addition, levels of p53 are tightly regulated by mdm2, which binds to p53 repressing its activity and triggering its degradation. The MDM2 gene is itself under the transcriptional control of p53, creating an autoregulatory feedback loop (2).

Alterations in the TP53 gene appear to be uncommon in prostate cancer, and their clinical significance has not been fully investigated. A recognized limitation of most studies is that they are confined to the analysis of p53 alterations, without analyzing other critical components that regulate its functions. The MDM2 gene is amplified in a variety of tumors, and mdm2 overexpression without amplification appears to be a common mechanism of p53 inactivation in certain cancers (3,4). Lack of data regarding the functional status of the p53 products encountered in the tumors analyzed represents another drawback. It has been reported that p21/WAF1 gene expression may serve as an indicator of p53 activity, since p21/WAF1 is under the transcriptional control of p53. However, serum or individual growth factors, such as epidermal growth factor (EGF), and fibroblast growth factor (FGF), were shown to induce p21 expression in p53-deficient cells (5,6). Thus, there are at least two separate pathways accounting for the induction of p21, one linked to DNA-damage recognition, and the other produced by signaling mechanisms caused by certain cellular mitogens.

In the present study, we have analyzed the patterns of p53 expression and those of critical components of its pathway, namely mdm2 and p21, in 86 patients with prostate cancer.

The association between these markers and clinicopathological parameters of poor outcome, including time to PSA relapse and proliferative index, were also examined.

5

#### EXPERIMENTAL DETAILS

##### Experimental Details for Second Series of Experiments

#### MATERIALS AND METHODS

10 Patients. A total number of 86 patients who underwent radical prostatectomy at Memorial Sloan-Kettering Cancer Center in the period between 1990 through 1991 were studied. Patient selection was based on the availability  
15 of both adequate clinical follow up and representative archival pathological materials for immunohistochemical analysis. The median age at the time of surgery was 65 years (range 46-74). Their median follow up was 64.5 months (range 10-94 months). Formalin-fixed, paraffin  
20 embedded prostate tissues were obtained from our archival tumor bank. Representative hematoxylin-eosin stained sections were examined to evaluate the histopathological characteristics of each case.

25 Clinicopathological parameters examined include pre-treatment PSA, pathologic stage and Gleason score, both determined based on the radical prostatectomy specimen. Time to PSA relapse was calculated from the day of surgery to the first detectable PSA. PSA relapse was defined as  
30 three consecutive rise in PSA at least one week apart. Only patients who had undetectable PSA level after surgery were included in this analysis.

Tumors were staged pT2 (n=51) and pT3 (n=35). Twenty-nine  
35 patients were Gleason score <7, while 18 patients were



Gleason  $\geq 7$ . In six cases, due to scarcity of tumor representation in the specimen, grade was considered to be not interpretable. Thirty-three patients (38.3%) received neoadjuvant hormone treatment preoperatively, and were defined as hormone-treated. These patients had non-evaluable Gleason scores. Patients who did not receive neoadjuvant hormone treatment were defined as hormone-naïve.

Monoclonal Antibodies and Immunohistochemistry. The following well characterized mouse monoclonal antibodies and corresponding final working dilutions were used for the present study: anti-p53 monoclonal antibody PAB1801 (Ab-2 clone; CalBiochem/Oncogene Science, Boston, MA; 1:500 dilution); anti-mdm2 monoclonal antibody 2A10 (a gift from Dr. Arnold Levine, Rockefeller University, New York, N.Y.; 1:500 dilution); and an anti-p21 monoclonal antibody (Ab-1 clone; CalBiochem/Oncogene Science; 1:20 dilution). An anti-Ki67 mouse monoclonal antibody (clone MIB1; Immunotech SA, France; 1:50 dilution) was used to assess proliferative index. MlgS-Kp1, a mouse monoclonal antibody of the same subclass as the primary antibodies listed above was used as negative control.

An avidin-biotin immunoperoxidase method was utilized. Briefly, sections were subsequently immersed in boiling 0.01% citric acid (pH 6.0) for 15 minutes to enhance antigen retrieval and incubated with primary antibodies overnight at 4°C. Biotinylated horse anti-mouse IgG antibodies were applied for 1h (Vector Laboratories, Burlingame, CA; 1:500 dilution), followed by avidin-biotin peroxidase complexes for 30 minutes (Vector Laboratories; 1:25 dilution). Diaminobenzidine was used as the final chromogen and hematoxylin was used as the nuclear counterstain. Nuclear immunoreactivity were classified on

a continuous scale with values that ranged from undetectable levels or 0% to homogeneous staining or 100%.

Statistical Analysis. The three markers were analyzed both  
5 as percentage of tumor cells and as discrete variables  
based on *a priori* cut-points. The cut-point for p53 of  
>20% was based on our previous analysis of p53 alterations  
in bladder cancer that revealed a strong association  
10 between p53 point mutation and p53 nuclear accumulation in  
>20% of tumor cells (7,8). For mdm2, the cut-point was  
based on what have been published correlating mdm2  
overexpression in  $\geq 20\%$  of tumor cells with worse  
clinicopathological parameters (9,10). The same principle  
15 applied to the Ki67 cut-point determination (11,12). For  
p21 the cut-point of >5% was based on our finding that  
normal prostate glands lack p21 expression, and the  
observation of p21 nuclear staining and presence of mitotic  
figures indicating high proliferative activity of the  
tumors.  
20  
The association of percentage of tumor cells expressing the  
markers with time to PSA relapse, while adjusting for other  
variables with known prognostic significance, was assessed  
using the Cox proportional hazards model (13). In  
25 addition, Kaplan-Meier estimation (14) was performed and  
the log rank test (15) employed to assess the univariate  
relationship between the individual markers using cut  
points and time to PSA relapse.  
30  
The associations between Gleason group and the three  
biomarkers were assessed using Fisher's exact test (16).  
Also, associations between the three markers and variables  
such as Ki67 proliferative index, stage, and hormone status  
were also assessed using the above test.  
35

# EXPERIMENTAL RESULTS

## Experimental Results for the Second Series of Experiments

5 Table 1 summarizes the data in relation to  
 clinicopathological parameters, including pre-treatment  
 PSA, tumor stage, Gleason tumor grade, hormone status,  
 proliferative index, and immunophenotype profile. Figure  
 7 illustrates the univariate relationships of the three  
 10 markers with time to PSA relapse with Kaplan-Meier curves  
 estimated.

Table 1. Summary Of Data In Relation To Immunophenotype Profile

|                                 | P53  |      | p21   |      | mdm-2 |      |
|---------------------------------|------|------|-------|------|-------|------|
|                                 | (#)  | (%)  | (#)   | (%)  | (#)   | (%)  |
| <u>PSA</u>                      |      |      |       |      |       |      |
| <4                              | 0/18 | 0    | 5/18  | 27   | 3/18  | 16   |
| 4-10                            | 3/28 | 10   | 7/28  | 25   | 7/28  | 25   |
| >10                             | 3/40 | 7.5  | 16/40 | 40   | 18/40 | 45   |
| p value                         |      | .374 |       | .382 |       | .060 |
| <u>Stage</u>                    |      |      |       |      |       |      |
| T<3                             | 3/51 | 5.8  | 14/51 | 27   | 11/51 | 21   |
| T=3                             | 3/35 | 8.5  | 14/35 | 40   | 17/35 | 48   |
| p value                         |      | .631 |       | .222 |       | .009 |
| <u>Gleason Score</u>            |      |      |       |      |       |      |
| <7                              | 0/29 | 0    | 7/29  | 24   | 7/29  | 24   |
| =7                              | 2/18 | 11   | 10/18 | 22   | 9/18  | 50   |
| NE                              | 4/33 | 2    | 10/33 | 30   | 11/33 | 33   |
| p value                         |      | .157 |       | .074 |       | .190 |
| <u>Hormone Status</u>           |      |      |       |      |       |      |
| Naive                           | 2/53 | 3    | 18/53 | 33   | 17/53 | 32   |
| Treated                         | 4/33 | 12   | 10/33 | 30   | 11/33 | 33   |
| p value                         |      | .139 |       | .725 |       | .904 |
| <u>Proliferation Index Ki67</u> |      |      |       |      |       |      |
| Low                             | 5/75 | 6    | 20/75 | 26   | 22/75 | 29   |
| High                            | 1/11 | 9    | 8/11  | 72   | 6/11  | 54   |
| p value                         |      | .768 |       | .002 |       | .096 |

p53 nuclear overexpression of >20% was observed in 6 of 86 cases. The distribution of p53% expression was primarily patients expressing less than 5% p53 (n=76). The other 10

patients had varying levels of p53% expression, indicating a very low frequency of p53 alteration in this group of patients. There is no correlation between p53 positive phenotype and pretreatment PSA, tumor stage, tumor grade, hormone status, or high proliferative index. Also, there is no association between p53 overexpression and p21 or mdm2 overexpression. A significant association was observed between p53 status determined by the cut-point and time to PSA relapse. This association is illustrated in Figure 7. Using the log rank test to examine the overall differences between p53 negative phenotype and p53 positive phenotype revealed a statistical significant difference  $P < 0.01$ . This indicates an obvious PSA relapse time advantage for patients who do not overexpress p53. However, the magnitude of this difference may not be reliably estimated due to the small number of patients and events in the p53 positive phenotype group.

mdm2 nuclear overexpression of  $\geq 20\%$  tumor cells was observed in 28 of 86 cases (32.5%). mdm2 positive phenotype was associated with advanced stage ( $P = 0.009$ ). In addition, mdm2 overexpression was observed not to be significant with respect to a decreased time to PSA relapse (Figure 7). A trend was observed between mdm2 overexpression and higher pretreatment PSA ( $P = 0.06$ ).

p21 nuclear overexpression of  $> 5\%$  tumor cells was observed in 28 of 86 patients (32.5%). Patients with p21 positive phenotype were observed to have a significant association with high Ki67 proliferative index ( $P = 0.002$ ). High Ki67 proliferative index was identified in 11 of 86 patients (12.7%). Patients with p21 positive phenotype had a significant association with decreased time to PSA relapse, as illustrated in Figure 7. Also, p21 overexpression was associated with mdm2 overexpression ( $P < 0.01$ ). However, no

association was observed between identification of p21 and/or mdm2 positive phenotype and p53 overexpression.

Forty-three of the total 86 patients had one or more altered markers. Patients with any alteration (p53 or mdm2 or p21) were observed to have a higher rate of PSA relapse ( $P < 0.01$ ).

The multivariate relationship between the markers and time to PSA relapse was assessed using Cox proportional hazards model. It was of interest to examine the effect of the markers while adjusting for variables with known prognostic significance. Both, p53 and p21 positive phenotypes were significant while adjusting for pre-treatment PSA and Gleason group ( $P < 0.01$  for both markers). Examination of the overexpression of at least one marker (p53, mdm2 or p21) with respect to time to PSA relapse showed that this variable was also significant ( $P < 0.01$ ) while adjusted for pre-treatment PSA and Gleason group. Tumor stage ( $< 3$  vs.  $\geq 3$ ) was not significant in either the univariate or multivariate analyses, and was thus excluded from the model. The model that seemed to account for the most information included p53 and p21, along with pre-treatment PSA.

## EXPERIMENTAL DISCUSSION

### Experimental Discussion for the Second Series of Experiments

Reports dealing with the frequency of TP53 mutations and p53 overexpression in prostate cancer have yielded conflicting results, alterations ranging from 2% to 65% of cases studied (17-21). This discrepancy might be explained

by the relatively small number of cases and different disease stages analyzed in some reports, the distinct methodologies employed, and the cutoff points used for evaluation of IHC results. However, a general finding was the association between p53 alterations and clinicopathological parameters of poor clinical outcome, such as high grade and late stage (18,19,22). In this study, we observed a relatively low frequency of p53 nuclear overexpression in patients with localized prostate cancer, as previously reported (23,24). To determine the potential clinical relevance of identifying a p53 positive phenotype, we correlated phenotypic characteristics of the tumors with the time to PSA relapse. This is considered the most sensitive indicator of success or failure following radical prostatectomy in patients treated for localized disease. Analysis of data revealed that p53 overexpression was significantly associated with PSA relapse ( $P < 0.01$ ) and independent of pretreatment PSA and Gleason group. However, the magnitude of this difference may not be reliably estimated due to the small number of patients and events in the positive phenotype. We also observed that all patients who received neoadjuvant hormone treatment prior to surgery and had tumors that overexpressed p53 relapsed. This finding could be due to the advanced stage at which patients presented and were selected for treatment using this modality. Mechanistically, an altered p53 status in this setting could have conferred resistance to castration-induced apoptosis, ultimately leading to disease relapse. The association between p53 overexpression and hormone refractory prostate cancer has been reported in locally advanced and metastatic disease (25). However, to our knowledge, this is the first report to suggest that this association might be an early event in the evolution of hormone refractory disease in clinically localized prostate

cancer.

In the present study we also analyzed alterations affecting other regulators of the p53 pathway in primary prostate cancer, including mdm2 and p21. The MDM2 gene maps to 12q13 and is found overexpressed in certain tumors, due to its amplification as a component of an amplicon that includes other relevant genes, such as CDK4. The MDM2 is under transcriptional regulation by p53, and encodes a 90-kDa zinc finger protein (mdm2) which contains a p53-binding site (26). It has been shown that mdm2 binds to p53, and acts as a negative regulator by inhibiting p53 transcriptional activity and targeting its degradation, thus creating an autoregulatory feedback loop (27). In this study, nuclear mdm2 overexpression was found in 32.5% of cases. We observed that mdm2 positive phenotype was significantly associated with advanced stage. It has been previously reported that MDM2 is not amplified on primary prostate cancer, based on a study of 29 tumors analyzed by Southern blot hybridization (28). The discrepancy between the rate of MDM2 gene amplification and protein overexpression has been described in Burkitt's lymphoma and breast cancer (29,30). Furthermore, it was observed in soft tissue sarcomas that mdm2 overexpression, rather than its amplification, was associated with worse clinical outcome (10). Based on data from this study, we can postulate that mdm2 overexpression is a frequent mechanism of p53 inactivation in prostate cancer, and in this context the MDM2 gene can be classified as an oncogene in this setting.

The p21/WAF1 gene encodes a nuclear protein member of the cyclin-dependent kinase inhibitory KIP family involved in senescence and cell quiescence (31). The p21/WAF1 gene is also transcriptionally regulated by p53. However, p21

induction could also be accomplished by a p53-independent pathway. Serum or individual growth factors, such as EGF and FGF, were shown to induce p21 in p53-deficient cells (32). Based on these data, it has been postulated that p21 induction could be activated through two separate pathways. The rate of p21/WAF1 mutations in human cancer is very low (33). However, there is an association between altered patterns of p21 expression and clinical outcome in certain tumors, such as bladder, colon, and hepatocellular carcinomas (34-36). Lack of p21 expression in these studies was correlated with poor clinical outcome, an expected finding if one postulates that p21 deficiency reflects p53 inactivation. As a corollary to this hypothesis, the p21 negative phenotype observed in the above referred studies was usually associated with p53 alterations. However, in our study we found that p21 positive phenotype was significantly associated with high proliferative index and mdm2 overexpression, but not with p53 status. Moreover, patients with p21 positive phenotype had a significant association with decreased time to PSA relapse. p21 overexpression has been reported to be associated with worse prognosis in other tumor types, including, breast, esophageal carcinoma, and squamous cell carcinomas of head and neck (37-39). Moreover, p21 overexpression was found to be associated with resistance to chemotherapy in acute myeloid leukemia and glioblastoma (40,41).

These data could be interpreted as follows (see Figure 8). A positive p21 phenotype could signify activation of p53 in response to DNA damage or cellular stress. This effect would result in G1 arrest of the prostate tumor cells expressing p21. We observed, on the contrary, an association between p21 positive phenotype and increased proliferative activity. Thus, it is more plausible to



postulate that the p21 overexpression observed is caused by a p53-independent transactivation mechanism. In the setting of prostate cancer, the alternative mechanism could be due to mitogenic stimuli via growth factor signaling.

5 There is abundant evidence regarding the upregulation of growth factor receptor/ligand activity in prostate tumors (42-46). An additional aberration causing p53 inactivation would be required in this model to explain the lack of cell death and association with proliferative activity. It is

10 our hypothesis that the increased mdm2 expression discussed above provides this requirement, further supporting the oncogenic role of mdm2 in prostate cancer.

Finally, the association between p21 and high proliferative

15 index might also reflect deregulated cyclinD1/CDK4 activity. In fact, we observed a strong association between p21 positive phenotype and cyclin D1 overexpression in this cohort of patients (Drobnjak et al, personal communication). Taken together, these data supports the

20 concept that p21 overexpression denotes an inefficient pRB control on S-phase entry.

Growth control in mammalian cells is accomplished largely by the action of the RB protein, regulating exit from the

25 G1 phase, and the p53 protein, triggering growth arrest or apoptotic processes. In this group of patients, there is enough evidence to suggest that both mechanisms are defective in prostate cancer. The high proliferative index reflects the inefficient pRB control. We postulate that

30 this phenomenon is produced by deregulated cyclinD1/CDK4 activity, which is associated with a p21 positive phenotype. The deactivation of a p53-dependent apoptosis could be explained by the degradation of p53 induced by mdm2 overexpression.

35

In sum, alterations affecting the p53 pathway are frequent events in prostate cancer. It is our hypothesis that a pathway of prostate cancer progression involves p53 inactivation caused by mdm2 overexpression, and that p21  
5 transactivation in this setting is due to an alternative signaling system rather than through a p53-dependent mechanism.

REFERENCES FOR SECOND SERIES OF EXPERIMENTS

1. Cordon-Cardo, C. Mutation of cell cycle regulators. Biological and clinical implication for human neoplasms. Am J Pathol, 147: 545-560,1995.
2. Momand, J., Zambetti, G.P., Olsen, D.C., George, D.L., and Levine A.J. The MDM2 oncogene products forms a complex with the p53 protein and inhibits p53-mediated transactivation. Cell, 69:1237-1245,1992.
3. Lianes, P., Orlow, I., Zhang, Z-F., Oliva, M., Sarkis, A., Reuter, V., and Cordon-Cardo., C. Altered patterns of MDM2 and TP53 expression in human bladder cancer. J Natl. Cancer Inst., 86:1325-1330, 1994.
4. Gorgoulis, V.G., Rassidakis,G.Z., Karameris, A.M., Papastamatiou, H., Trigidou, R., Veslemes, M.,Rassidakis, A.N., and Kittas, C. Immunohistochemical and molecular evaluation of the mdm2 gene product in bronchogenic carcinoma. Mod Pathol,9:544-554,1996.
5. Sinicrope, F. A., Roddey, G., Lemoine, M., Ruan, S., Stephens, L. C., Frazier, M.L., Shen, Y., and Zhang,W. Loss of p21WAF1/Cip1 protein expression accompanies progression of sporadic colorectal neoplasms but not hereditary nonpolyposis colorectal cancers. Clin Cancer Res, 4:1251-1261,1998.
6. Lacombe, L., Orlow, I., Zhang, Z-F, Oliva, M., Sarkis, A., Reuter, V. G, and Cordon-Cardo,C. Analysis of p21WAF1/CIP1 in primary bladder tumors. Ocol Res, 8:409-414,1996.

7. Sarkis, A.S., Dalbagni, G., Cordon-Cardo, C., Zhang, Z-F., Sheinfeld, J., Fair, W.R., Herr, H.W., and Reuter, V.E. Nuclear overexpression of p53 protein in transitional cell bladder carcinoma: a marker for disease progression. J Natl. Cancer Inst., 85:53-59, 1993.
8. Cordon-Cardo, C., Dalbagni, G., Saez, G.T., Oliva, M.R., Zhang, Z-F., Rosai, J., Reuter, V.E., and Pellicer, A. p53 mutations in human bladder cancer: genotypic versus phenotypic patterns. Int J. Cancer, 56:347-353, 1992.
9. Osman, I., Scher, H.I., Zhang, Z-F., Pellicer, I., Hamza, R., Eissa, S., Khaled, H., and Cordon-Cardo, C. Alterations affecting the p53 control pathway in Bilharzial-related bladder cancer. Clin Cancer Res, 3:531-536, 1997.
10. Cordon-Cardo, C., Latres, E., Drobnjak, M., Oliva, M.R., Pollack, D., Woodruff, J.M, Marechal, V., Chen, J., Brennan, M.F., and Levine, A.J. Molecular abnormalities of mdm2 and p53 genes in adult soft tissue sarcomas. Cancer Res, 54:794-799, 1994.
11. Heslin, M.J., Cordon-Cardo, C., Lewis, J.J., Woodruff, J.M., and Brennan, M.F. Ki67 detected by MIB-1 predicts distant metastasis and tumor mortality in primary, high grade extremity soft tissue sarcoma. Cancer, 83 :490-497, 1998.
12. Osman, I., Scher, H., Zhang, Z-F., Soos, T.J., Hamza, R., Eissa, S., Khaled, H., Koff, A., and Cordon-Cardo, C. Expression of cyclin D1, but not cyclins E and A, is related to progression in

Bilharzial bladder cancer. Clin Cancer Res, 3:2247-2251, 1997.

- 5 13. Cox, D.R. Regression models and life tables. J Royal Statistical Soc, 34:187-220, 1972.
14. Kaplan, E. L. and Meier, P. Nonparametric estimation from incomplete observations. J American Statistical Assoc, 53:457-481, 1958.
- 10 15. Mantel, N. Evaluation of survival data and two new rank order statistics arising in its consideration. Cancer Chemo Reports, 50:163-170, 1966.
- 15 16. Agresti, A. Categorical data Analysis. pp39-44. Wiley, New York 1996.
17. Hall, M.C, Navone, N.M., Troncoso, P., Pollack, A., Zagars, G.K., von Eschenbach, A.C., Conti, C.J., and Chung, L.W. Frequency and characterization of p53 mutations in clinically localized prostate cancer. Urology, 45:470-475, 1995.
- 20 18. Kubota, Y., Shuin, T., Uemura, H., Fujinami, K., Miyamoto, H., Torigoe, S., Dobashi, Y., Kitamura, H., Iwasaki, Y., and Danenberg, K. Tumor suppressor gene p53 mutations in human prostate cancer. Prostate, 27:18-24, 1995.
- 25 19. Hughes, J.H., Cohen, M.B., and Robinson, R.A. p53 immunoreactivity in primary and metastatic prostatic adenocarcinoma. Mod Pathol, 8:462-466, 1995.
- 30 20. Theodorescu, D., Broder, S.R., Boyd, J.C., Mills, S.E., and Frierson, H.F Jr. p53, bcl-2 and
- 35

retinoblastoma proteins as long-term prognostic markers in localized carcinoma of the prostate. J Urol, 158:131-137,1997.

- 5      21. Gumerlock, P.H., Chi, S.G., Shi, X.B., Voeller, H.J., Jacobson, J.W., Gelmann, E.P., and deVere White, R.W. p53 abnormalities in primary prostate cancer: single-strand conformation polymorphism analysis of complementary DNA in comparison with  
10      genomic DNA. The Cooperative Prostate Network. J Natl Cancer Inst., 89:66-71, 1997
- 15      22. Bauer, J.J., Sesterhenn, I.A., Mostofi, F.K., McLeod, D.G., Srivastava, S., and Moul, J.W. Elevated levels of apoptosis regulator proteins p53 and bcl-2 are independent prognostic biomarker in surgically treated clinically localized prostate cancer. J Urol , 156:1511-1516, 1996.
- 20      23. Sinik, Z., Alkibay, T., Ataoglu, O., Biri, H., Sozen, S., Deniz, N., Karaoglan, U., and Bozkirli, I. Nuclear p53 overexpression in bladder, prostate, and renal carcinomas. Int J Urol,4:546-551,1997
- 25      24. Brooks, J.D., Bova, G.S., Ewing, C.M., Piantadosi, S., Carter, B.S., Robinson, J.C., Epstein, J.I, and Isaacs, W.B. An uncertain role for p53 gene alterations in human prostate cancers. Cancer Res, 56:3814-3822, 1996.
- 30      25. Srivastava, S., and Moul, J. W. p53 tumor suppressor gene alteration in prostate cancer and potential gene therapy approaches. Molecular Urolog, 1:151-158, 1997.

26. Kubbutat, M.H.G., Ludwig, R.L., Ashcroft, M.,  
Vousden, K.H. Regulation of Mdm2-directed  
degradation by the C terminus of p53. *Mol Cell Biol*,  
18:5690-5698, 1998.
- 5
27. Grossman, S.R., Perez, M., Kung, A.L., Joseph, M.,  
Mansur, C., Xiao, Z.X., Kumar, S., Howley, P..M.,  
and Livingston, D.M. p300/MDM2 complexes  
participate in MDM2-mediated p53 degradation. *Mol*  
10 *Cell*, 2:405-415, 1998.
28. Ittmann, M., Wieczorek, R., Heller, P., Dave, A.,  
Provet, J., and Krolewski, J. Alterations in the  
p53 and MDM-2 genes are infrequent in clinically  
15 localized, stage B prostate adenocarcinomas. *Am J*  
*Pathol*, 145:287-293, 1994.
29. Capoulade, C., Bressac-de Paillerets, B., Lefrere,  
I., Ronsin, M., Feunteun, J., Tursz, T., Wiels,  
20 Overexpression of MDM2, due to enhanced translation,  
results in inactivation of wild-type p53 in Burkitt's  
lymphoma cells. *Oncogene*, 16:1603-1610, 1998.
30. Bueso-Ramos, C.E., Manshour, T., Haidar, M.A.,  
25 Yang, Y., McCown, P., Ordonez, N., Glassman, A.,  
Sneige, N., and Albitar, M. Abnormal expression of  
MDM-2 in breast carcinomas. *Breast Cancer Res Treat*,  
37:179-188, 1996.
- 30 31. Kuzumaki, T., Kobayashi, T., and Ishikawa, K.  
Genistein induces p21(Cip1/WAF1) expression and blocks  
the G1 to S phase transition in mouse fibroblast and  
melanoma cells. *Biochem Biophys Res Commun* ,  
251:291-295, 1998.

32. Fan, Z., Shang, B.Y., Lu, Y., Chou, J.L., and Mendelsohn, J. Reciprocal changes in p27(Kip1) and p21(Cip1) in growth inhibition mediated by blockade or overstimulation of epidermal growth Factor receptors. Clin Cancer Res, 3:1943-1948,1997.
33. Elbendary, A.A., Cirisano, F.D., Evans, A.C Jr., Davis, P.L., Iglehart, J.D., Marks, J.R., and Berchuck, A. Relationship between p21 expression and mutation of the p53 tumor suppressor gene in normal and malignant ovarian epithelial cells. Clin Cancer Res, 2:1571-1575,1996.
34. Stein, J.P., Ginsberg, D.A., Grossfeld, G.D., Chatterjee, S.J., Esrig, D., Dickinson, M.G., Groshen, S., Taylor, C.R., Jones, P.A., Skinner, D.G., and Cote, R.J. Effect of p21WAF1/CIP1 expression on tumor progression in bladder cancer. J Natl. Cancer Inst., 90:1072-1079, 1998.
35. Valassiadou, K.E., Stefanaki, K., Tzardi, M., Datseris, G., Georgoulas, V., Melissas, J., Tsiftsis, D.D., Delides, G., Kanavaros, P. Immunohistochemical expression of p53, bcl-2, mdm2 and waf1/p21 proteins in colorectal adenocarcinomas. Anticancer Res, 17:2571-2576,1997.
36. Qin, L.F., Ng, I.O., Fan, S.T., and Ng, M. p21/WAF1, p53 and PCNA expression and p53 mutation status in hepatocellular carcinoma. Int J Cancer, 79:424-428, 1998.
37. Caffo, O., Doglioni, C., Veronese, S., Bonzanini, M., Marchetti, A., Buttitta, F., Fina, P., Leek, R., Morelli, L., Palma, P.D., Harris, A.L., and



Barbareschi, M. Prognostic Value of p21(WAF1) and p53 Expression in Breast Carcinoma: An Immunohistochemical Study in 261 Patients with long-term Follow-Up. Clin Cancer Res, 2:1591-1599,1996.

5

38. Sarbia, M., Stahl, M., Hausen, A. Z., Zimmermann, K., Wang, L., Fink, U., Heep, H., Dutkowski, P., Willers, R., Muller, W., Seeber, S., and Gabbert, H. Expression of p21 predicts outcome of esophageal cancer patients treated by surgery alone or by combination therapy modalities. Clin Cancer Res, 4:2615-2623,1998.

10

39. Erber, R., Klein, W., Andl, T., Enders, C., Born, A.I., Conradt, C., Bartek, J., and Bosch, F.X. Aberrant p21(CIP1/WAF1) protein accumulation in head-and-neck cancer. Int J Cancer, 74:383-389,1997.

15

40. Zhang, W., Kornblau, S.M., Kobayashi, T., Gambel, A., Claxton, D., and Deisseroth, A.B. High levels of constitutive WAF1/Cip1 protein are associated with chemoresistance in acute myelogenous leukemia. Clin Cancer Res, 1:1051-1057,1995.

20

41. Ruan, S., Okcu, M.F., Ren, J.P., Chiao, P., Andreeff, M., Levin, V., and Zhang, W. Overexpressed WAF1/Cip1 renders glioblastoma cells resistant to chemotherapy agents 1,3-bis(2-chloroethyl)-1-nitrosourea and cisplatin. Cancer Res, 58:1538-1543,1998.

25

42. Tanaka, A., Furuya, A., Yamasaki, M., Hanai, N., Kuriki, K., Kamiakito, T., Kobayashi, Y., Yoshida, H., Koike, M., and Fukayama M. High frequency of fibroblast growth factor (FGF) 8 expression in clinical prostate cancers and breast tissues,

30

35

immunohistochemically demonstrated by a newly established neutralizing monoclonal antibody against FGF 8. Cancer Res, 58:2053-2056, 1998.

- 5      43. Peehl, D.M., and Sellers, R.G. Basic FGF, EGF, and PDGF modify TGFbeta-induction of smooth muscle cell phenotype in human prostatic stromal cells. Prostate, 35:125-134, 1998.
- 10      44. Culig, Z., Hobisch, A., Cronauer, M.V., Radmayr, C., Hittmair, A., Zhang, J., Thurnher, M., Bartsch, G., and Klocker, H. Regulation of prostatic growth and function by peptide growth factors. Prostate, 28:392-405, 1996.
- 15      45. Scher, H.I., Sarkis, A., Reuter, V., Cohen, D., Netto, G., Petrylak, D., Lianes, P., Fuks, Z., Mendelson, J., and Cordon-Cardo, C. Changing pattern of expression of the epidermal growth factor receptor and transforming growth factor alpha in the progression of prostatic neoplasms. Clin Cancer Res 1:545-550, 1995.
- 20      46. Cohen, D.W., Simak, R., Fair, W.R., Melamed, J., Scher, H.I., and Cordon-Cardo, C. Expression of transforming growth factor-alpha and the epidermal growth factor receptor in human prostate tissues. J Urol 152:2120-2124, 1994.
- 25

THIRD SERIES OF EXPERIMENTS

Cyclin D1 is a key regulator of the G1 phase progression of the cell division cycle. There is an increasing evidence  
5 that deregulated cyclin D1 expression is implicated in tumorigenesis and tumor progression in certain neoplasms. The present study was conducted in order to analyze the alterations affecting cyclin D1 in prostate cancer, as well as to assess its potential clinical significance. We  
10 studied 116 cases of primary (n=86) and metastatic (n=30) prostate carcinomas using immunohistochemistry and a well characterized monoclonal antibody to cyclin D1. The results were correlated with proliferative index, as assessed by Ki67 antigen expression and with  
15 clinicopathologic variables of poor prognosis. Cyclin D1 positive phenotype, defined as identification of immunoreactivity in the nuclei of  $\geq 20\%$  tumor cells, was found in 26 of 116 (22%) cases. A significant association was observed between cyclin D1 positive phenotype and  
20 clinicopathologic parameters, such as advanced tumor stage ( $T \geq 3$ ) ( $P=0.045$ ), evidence of bone metastases ( $P=0.001$ ) and with elevated preoperative prostate specific antigen measurements (PSA  $>10$  ng/ml) ( $P=0.01$ ). Ki67 proliferative index was considered high when  $\geq 20\%$  tumor cells displayed  
25 positive nuclear staining, a phenotype that was observed in 20 of 107 (19%) evaluable cases. Moreover, high Ki67 proliferative index was associated with cyclin D1 overexpression ( $P=0.01$ ). These data support the hypothesis that alterations of cyclin D1 may represent an oncogenic  
30 event in human prostate cancer. Furthermore, it appears that cyclin D1 overexpression contributes to tumor progression in a subset of particularly aggressive prostate carcinomas, especially those developing osseous metastases.

35 Prostate cancer has been reported to be a neoplastic

disease of a slow growth rate. Nevertheless, it still represents the second leading cause of cancer deaths in men in the United States. There is an obvious discrepancy between the clinical impression of a slowly growing neoplasm and tendency to produce an aggressive metastatic disease in individual patients. The prognostic indicators of histologic grade, pathologic stage, DNA ploidy, and tumor cell proliferative index proved to be of limited value in determining the biologic behavior of prostate cancer (1-3). Tumor suppressor genes, particularly p53 and RB, implicated in the molecular genetics of many human malignancies, were reported to be altered in a rather low frequency in prostate cancer (4-8).

Cell cycle transitions are controlled by functional heterodimers composed of a cyclin, acting as a regulatory subunit, and cyclin-dependent kinase (Cdk), which acts as the catalytic component (9). Multiple cyclins have been isolated and characterized, and a temporal map of their expression has been delineated. It is postulated that the complexes formed by cyclin D1 and Cdk4 govern G1 progression, while cyclin E-Cdk2 controls entry into S-phase and cyclin A-Cdk2 affects the regulation through S-phase (10). Cyclin D1-Cdk4 complexes exert their function through the phosphorylation of the product encoded by the retinoblastoma gene, pRb, in order to overcome the cell cycle block imposed by hypophosphorylated pRb (10). Several studies suggest that gene amplification and overexpression of cyclin D1 and Cdk4 are oncogenic events in certain tumors, including breast cancer (11), head and neck tumors (12, 13), esophageal (14, 15) and colorectal carcinoma (16). We undertook this study in order to analyze patterns of cyclin D1 expression in prostate cancer. The alterations identified were correlated with Ki67 proliferative index, as well as relevant

clinicopathologic parameters, in an attempt to define their potential biologic significance in prostate cancer.

## EXPERIMENTAL DETAILS

5

### Experimental Details for Third Series of Experiments

## MATERIALS AND METHODS

10 Patients Characteristics and Tissues. A cohort of 116  
patients with prostate carcinoma were evaluated, consisting  
of 86 primary and 30 metastatic cases (eight metastases to  
lymph node and 22 metastases to bone). All primary tumors  
(n=86) represented consecutive cases of patients who  
15 underwent radical prostatectomy at Memorial Sloan-Kettering  
Cancer Center, in the period of 1990 and 1991. All  
metastatic cases (n=30) were selected on the basis of the  
availability of tissue in the tumor bank. Samples were  
formalin-fixed, paraffin embedded tissue specimens,  
20 obtained from the Department of Pathology at Memorial  
Sloan-Kettering Cancer Center. Representative hematoxylin-  
eosin stained sections of each paraffin block were examined  
microscopically to confirm the presence of tumor, as well  
as to evaluate the pathologic grade and stage of the tumors  
25 analyzed. Thirty-three of 86 patients with primary  
carcinoma received preoperatively neoadjuvant hormone  
therapy (hormone treated), while the remaining 53 patients  
were not treated with such protocols and were considered  
hormone-naive. Hormone-naive primary tumors with sufficient  
30 tumor representation on tissue sections were assigned  
histologic grade (n=47). Histologic grade was categorized  
into two groups: low grade (Gleason score <7), and high  
grade (Gleason score ≥7). According to pathologic stage,  
cases were grouped into early (organ confined tumors, T<sub>2</sub>),  
35 or advanced tumors (extending beyond prostatic capsule,

≥T<sub>3</sub>). The response variable time to prostate-specific antigen (PSA) relapse was defined as the time from radical prostatectomy to the time of the first detectable (non zero) PSA measurement. Three consecutive increases of PSA  
5 were required to confirm PSA relapse. Only patients who had a nonmeasurable PSA after radical prostatectomy were included in the analysis.

Monoclonal Antibodies and Immunohistochemistry. The  
10 following well characterized antibodies and corresponding final working concentrations were used for the present study: anti-cyclin D1 mouse monoclonal antibody (Ab-3, clone DCS-6, IgG1, Oncogene, Calbiochem, Cambridge, MA; 1μg/ml); anti-Ki67 mouse monoclonal antibody (clone MIB-1,  
15 IgG1, Immunotech, Marseille, France; 4μg/ml). A nonspecific mouse IgG1 kappa monoclonal antibody was used as a negative control at similar working concentrations. Immunohistochemistry was performed on 5μm tissue sections using avidin-biotin-peroxidase method and antigen  
20 retrieval. Briefly, sections were immersed in boiling 0.01 M citric acid (pH 6.0) and heated in microwave oven for 15 minutes, to enhance epitope exposure. After cooling to room temperature, slides were incubated with 10% normal horse serum for 30 minutes. Subsequently, appropriately  
25 diluted primary antibodies were applied for overnight incubation at 4°C. Biotinylated horse anti-mouse IgG antibodies were used as secondary reagents, applied for an incubation period of 30 minutes (Vector Laboratories, Burlingame, CA; 1:500 dilution), followed by avidin-biotin-peroxidase complexes incubated for 30 minutes (Vector  
30 Laboratories - 1:25 dilution). Diaminobenzidine was used as the final chromogen and hematoxylin as the nuclear counterstain.

Immunohistochemistry Evaluation. Nuclear immunoreactivities for both cyclin D1 and Ki67 antigens, were classified into two categories defined as follows: negative (<20% tumor cells displaying nuclear immunostaining), and positive (≥20% tumor cells with nuclear immunostaining). The appropriateness of this cutoff point was validated graphically by using predicted survival time and looking at specific immunoreactivities as a continuum data in this group of patients. Ultimately, results were interpreted as defined above.

Statistical Methods. The baseline variables examined were PSA (ng/ml) at time of diagnosis (divided into three categories: <4, 4-10 and >10), Tumor grade (Gleason score) (divided into two mutually exclusive categories: <7 or ≥7), Pathologic stage T (2 or ≥3), and percent cyclin D1 and Ki67 expression. Statistical analyses were conducted to assess: 1) the correlation between immunophenotypic variables and clinicopathologic parameters such as: presentation status, tumor grade, pathologic stage, preoperative PSA, and hormonal status; 2) the correlation among immunophenotypic variables; 3) association between immunophenotypes and PSA relapse free survival. The Mantel-Haenszel chi-square test was used to assess the associations among the different variables and results were considered significant if the P value was <0.05. The FREQ procedure in SAS was used for this study (17). The associations between time to PSA relapse and the immunophenotypes were evaluated using the Log Rank test and Kaplan Meier estimates (18).

## EXPERIMENTAL RESULTS

### Experimental Results for the Third Series of Experiments

- 5 Table 2 summarizes immunohistochemical data in relation to clinicopathologic parameters. Figure 9 illustrates the immunohistochemical staining patterns of cyclin D1 in representative cases of primary tumors and bone metastases.

**Table 2.** Clinicopathologic Parameters in Relation to Cyclin D1 Immunoreactivity

| Parameter                   | cyclin D1- (<20%) |         | cyclin D1+ (≥20%) |        | Total | P-value |
|-----------------------------|-------------------|---------|-------------------|--------|-------|---------|
|                             | N                 | (%)     | N                 | (%)    |       |         |
| Total Patients              | 90                | (77.6)  | 26                | (22.4) | 116   |         |
| Presentation                |                   |         |                   |        |       |         |
| Primary tms                 | 76                | (88.4)  | 10                | (11.6) | 86    |         |
| LN metastases               | 7                 | (87.5)  | 1                 | (12.5) | 8     | NS      |
| Primary tms                 | 76                | (88.4)  | 10                | (11.6) | 86    |         |
| Bone metastases             | 7                 | (31.8)  | 15                | (68.2) | 22    | 0.001   |
| Ki67 proliferative index    |                   |         |                   |        |       |         |
| Low (<20%)                  | 71                | (81.6)  | 16                | (18.4) | 87    |         |
| High (≥20%)                 | 11                | (55.0)  | 9                 | (45.0) | 20    | 0.01    |
| Primary tumors              | 76                | (88.4)  | 10                | (11.6) | 86    |         |
| Tm. Grade (Gleason)*        |                   |         |                   |        |       |         |
| Low (<7)                    | 27                | (93.1)  | 2                 | (6.9)  | 29    |         |
| High (≥7)                   | 17                | (94.4)  | 1                 | (5.6)  | 18    | NS      |
| Path. Stage                 |                   |         |                   |        |       |         |
| Early (T <sub>2</sub> )     | 48                | (94.1)  | 3                 | (5.9)  | 51    |         |
| Advanced (≥T <sub>3</sub> ) | 28                | (80.0)  | 7                 | (20.0) | 35    | 0.045   |
| Hormonal status             |                   |         |                   |        |       |         |
| H. naive                    | 49                | (92.5)  | 4                 | (7.5)  | 53    |         |
| H. treated                  | 27                | (81.8)  | 6                 | (18.2) | 33    | NS      |
| Pretreatment PSA            |                   |         |                   |        |       |         |
| <4 ng/ml                    | 17                | (94.4)  | 1                 | (5.6)  | 18    |         |
| 4-10 ng/ml                  | 28                | (100.0) | 0                 | (0.0)  | 28    |         |
| >10 ng/ml                   | 31                | (77.5)  | 9                 | (22.5) | 40    | 0.01    |

\*Patients who had received neoadjuvant hormonal therapy (n=33) and tissue sections with inadequate tumor representation (n=6) were not assigned Gleason grade and were excluded from this analysis.



Cyclin D1 was expressed in  $\geq 20\%$  tumor cells in 26 of 116 (22%) evaluable cases, corresponding to 10 of 86 (12%) primary lesions and 16 of 30 (53%) metastases. There was a statistically significant association between cyclin D1 overexpression and the presence of bone metastases. We observed that 15 of 22 (68%) bone metastases lesions overexpressed cyclin D1, while only 10 of 86 (12%) primary tumors presented with this positive phenotype ( $p=0.001$ ). Cyclin D1 overexpression was also associated with advanced pathologic stage in primary tumors. We found that 7 of 35 (20%) tumors of advanced stage (extending beyond the prostatic capsule,  $\geq T_3$ ) displayed cyclin D1 nuclear overexpression, compared to only 3 of 51 (6%) organ confined tumors ( $T_2$ ) ( $P=0.045$ ). Cyclin D1 was also detected at increased percentage of tumor cells in patients with high initial pretreatment PSA values. Nine of 68 (13%) patients with PSA  $\geq 10\text{ng/ml}$  were cyclin D1 positive compared to only 1 of 18 (5%) patients with PSA  $< 4\text{ng/ml}$  ( $P=0.01$ ). There was no association between cyclin D1 overexpression and tumor grade (Gleason score) or hormonal status (hormone naive vs. hormone treated). In order to assess disease progression we evaluated the time to PSA failure after radical prostatectomy. There was no association between cyclin D1 nuclear overexpression and early relapse as defined by increased PSA measurements after radical prostatectomy in these group of patients ( $p=0.2$ ).

Cyclin D1 overexpression correlated well with high Ki67 proliferative index, which was scored as being high in 20 of 107 (19%) evaluable tumors. Nine of 20 (45%) tumors displaying high Ki67 proliferative index also possessed cyclin D1 nuclear overexpression, while only 16 of 87 (18%) cases with low Ki67 proliferative index overexpressed cyclin D1 ( $P=0.01$ ). Nevertheless, Ki67 proliferative index

alone was not associated with clinicopathologic parameters of poor outcome in this cohort of patients.

## EXPERIMENTAL DISCUSSION

5

### Experimental Discussion for the Third Series of Experiments

Autopsy records show that by the age of 80, approximately 60-70% of men around the world, have histologic evidence of prostatic carcinoma (19). Although this fact indicates generally slow growing nature of this malignancy, there are vast differences in the progression rate and development of clinically evident or metastatic disease in these patients during their lifetime. Prostate cancer progression tends to follow periprostatic and perivascular penetration, invasion along perineural spaces, pelvic lymph node metastases and particularly bone metastases (20). Almost one fourth of newly diagnosed cases presents with lymph node and/or osseous metastases and only one fourth of those survive five years (21). We were interested in analyzing the molecular events that might be responsible for the progression of prostate cancer from indolent to a life threatening, metastatic disease.

Some earlier reports on determining prostate tumor proliferation, measured by flow cytometric S-phase fraction showed a positive predictive value of this variable and prostate cancer progression (2). Tumors that demonstrate a higher proliferation rate are more likely to grow to and beyond prostatic capsule and to produce distant metastases. Recently, the cell division cycle regulatory mechanisms and their oncogenic role have become a major focus of cancer research. There is ever growing literature on cyclins and their associated kinases and their role in tumorigenesis (22). Particularly D-family cyclins were implicated in

specific human tumors. Bartkova et al. (23) report on a large group of various human malignancies, including carcinoma of the breast, uterus, colon, melanomas and soft tissue sarcomas, high proportion of which exhibit immunoreactivity for cyclin D1. By far the most frequent chromosomal abnormality that affects cyclin D1 in majority of tumors is DNA amplification that results in increased expression of the RNA transcripts and protein levels (24). In some tumor types, however, immunohistochemistry proved to be the most accurate technique in determining deregulated expression of cyclin D1 (11, 23). In this study we used immunohistochemistry to determine cyclin D1 expression in patients with prostate carcinoma. To our knowledge this first expression study on cyclin D1 in both primary tumors and bone metastases specimens. Only recently Kallakury et al (25) evaluated the expression of p34<sup>cdc2</sup> and cyclin D1 in patients with radical prostatectomy. Results were correlated with conventional markers of poor prognosis. The authors showed no association between cyclin D1 immunoreactivity and clinicopathologic parameters, such as tumor grade, pathologic stage, lymph node metastases and with disease free survival. Our data in this study, on the other hand, suggest the involvement of cyclin D1 in the progression of human prostate cancer. There was a remarkably significant difference in the levels of cyclin D1 expression between bone metastases and primary tumors. There was an association between cyclin D1 immunoreactivity and tumors with advanced pathologic stage. Cyclin D1 further correlated well with high Ki67 proliferative index. Taken together, these results support the theory that increased levels of cyclin D1 expression contribute to cell cycle imbalance with extremely shortened G1 phase and possibly with reduced cell requirements for growth factors to proliferate (26, 27). Subclones of tumor cells with

elevated cyclins expression may acquire uncontrolled growth advantage and contribute to tumor progression.

5 In this study we conclude that cyclin D1 may play an  
oncogenic role in prostate cancer. Our data indicate that  
cyclin D1 is involved in tumor progression, particularly in  
a development of bone metastases.. However, in order to  
determine the timeframe and prognostic value of this marker  
10 in prostate cancer, from the early onset to the evolution  
of metastatic disease, we intend to evaluate cyclin D1  
immunophenotypes in paired samples of primary tumors and  
metastatic sites from the same patients.

15

REFERENCES FOR THIRD SERIES OF EXPERIMENTS

1. Gleason, D.F. Histologic grading of prostate cancer: a perspective. Hum. Pathol., 23:273-279, 1992.  
5
2. Visakorpi, T., Kallionemi, O.-P., Paronen, I.Y.I., Isola, J.J., Heikkinen, A.I., and Koivula, T.A. Flow cytometric analysis of DNA ploidy and S-phase fraction from prostatic carcinomas: implications for prognosis and response to endocrine therapy. Br. J. Cancer, 64: 578-582, 1991.  
10
3. Bubendorf, L., Sauter, G., Moch, H., Schmid, H.-P., Gasser, T.C., Jordan, P., and Mihatsch, M.J., Ki67 labeling index: an independent predictor of progression in prostate cancer treated by radical prostatectomy. J. Pathol., 178:437-441, 1996.  
15
4. Isaacs, W.B., Carter, B.S., and Ewing, C.M. Wild-type p53 suppresses growth of human prostate cancer cells containing mutant p53 alleles. Cancer Res., 51:4716-4720, 1991.  
20
5. Bookstein, R., Rio, P., Madreperla, S.A., Hong, F., Allred, C., Grizzle, W.E., and Lee, W.-H. Promoter deletion and loss of retinoblastoma gene expression in human prostate carcinoma. Proc. Natl. Acad. Sci. USA, 87:7762-7766, 1990.  
25
6. Visakorpi, T., Kallionemi, O.-P., Heikkinen, A., Koivula, T., and Isola, J. Small subgroup of aggressive, highly proliferative prostatic carcinomas defined by p53 accumulation. J.  
30  
35

Natl. Cancer Inst., 84:883-887, 1992.

7. Bookstein, R., MacGrogan, D., Hilsenbeck, S.G., Sharkey, F., and Allred, C.D. p53 is mutated in  
5 a subset of advanced-stage prostate cancers. Cancer Res., 53:3369-3373, 1993.
8. Navone, N.M., Troncoso, P., Pisters, L.L., Goodrow, T.L., Palmer, J.L., Nichols, W.W., von  
10 Eschenbach, A.C., and Conti, C.J. p53 protein accumulation and gene mutation in the progression of human prostate carcinoma. J. Natl. Cancer Inst., 85:1657-1669, 1993.
- 15 9. Sherr, C.J. G1 phase progression: cycling on cue. Cell, 79:551-555, 1994.
10. Matsushime, H., Quelle, D.E., Shurtleff, S.A., Shibuya, M., Sherr, C.J., and Kato, J.-Y. D-  
20 type cyclin-dependent kinase activity in mammalian cells. Mol. Cell Biol., 14:2066-2076, 1994.
11. Gillett, C., Fantl, V., Smith, R., Fisher, C.,  
25 Bartek, J., Dickson, C., Barnes, D., and Peters, G. Amplification and overexpression of cyclin D1 in breast cancer detected by immunohistochemical staining. Cancer Res., 54:1812-1817, 1994.
- 30 12. Jares, P., Fenandez, P.L., Campo, E., Nadal, A., Bosch, F., Aiza, G., Nayach, I., Traserra, J., and Cardesa, A. PRAD-1/Cyclin D1 gene amplification correlates with messenger RNA  
35 overexpression and tumor progression in human

laryngeal carcinomas. Cancer Res., 54:4823-4827, 1994.

- 5 13. Michalides, R., van Veelen, N., Hart, A., Loftus, B., Wientjens, E., and Balm, A. Overexpression of cyclin D1 correlates with recurrence in a group of forty-seven operable squamous cell carcinomas of the head and neck. Cancer Res., 55:975-978, 1995.
- 10 14. Jiang, W., Kahn, S.M., Tomita, N., Zhang, Y.-J., Lu, S.-H., and Weinstein, B.I. Amplification and expression of the human cyclin D gene in esophageal cancer. Cancer Res., 52:2980-2983, 15 1992.
- 20 15. Naitoh, H., Shibata, J., Kawaguchi, A., Kodama, M., and Hattori, T. Overexpression and localization of cyclin D1 mRNA and antigen in esophageal cancer. Am. J. Pathol., 146:1161-1169, 1995.
- 25 16. Zhang, T., Nanney, L.B., Luongo, C., Lamps, L., Heppner, K.J., DuBois, R.N., and Beauchamp, R.D. Concurrent overexpression of cyclic D1 and cyclin-dependent kinase 4 (Cdk4) in intestinal adenomas from multiple intestinal neoplasia (Min) mice and human familial adenomatous polyposis patients. Cancer Res., 57:169-175, 30 1997.
17. SAS Institute Inc., SAS/STAT user guide, version 6 Cary NC. SAS Institute Inc., 1990.
- 35 18. Kaplan, E.L., and Meier, P. Nonparametric

estimation from incomplete observations. J. Am. Stat. Assoc., 53:457-481, 1958.

- 5 19. Carter, B.H., Piantadosi, S., and Isaacs, J.T. Clinical evidence for and implications of the multistep development of prostate cancer. J. Urol., 143:742-746, 1990.
- 10 20. Raghavan, D., Scher, H.I., Leibel, S.A., and Lange, P. Principles and practice of genitourinary oncology. Lippincott-Raven publishers. Philadelphia. New York, 1997.
- 15 21. Murphy, G.P., Mettlin, C., Menck, H., Winchester, D.P., and Davidson, A.M. The national survey of prostate cancer in the United States by the American College of Surgeons. J. Urol., 127:928-934, 1982.
- 20 22. Hall, M., and Peters, G. Genetic alterations of cyclins, cyclic-dependent kinases, and cdk inhibitors in human cancer. Adv. Cancer Res., 68:67-108, 1996.
- 25 23. Bartkova, J., Lukas, J., Strauss, M., and Bartek, J. Cyclic D1 oncoprotein aberrantly accumulates in malignancies of diverse histogenesis. Oncogene, 10:775-778, 1995.
- 30 24. Lammie, G.A., Fantl, V., Smith, R., Schuuring, E., Brookes, S., Michalides, R., Dickson, C., Arnold, A., and Peters, G. D11S287, a putative oncogene on chromosome 11q13, is amplified and expressed in squamous cell and mammary carcinomas and linked to BCL-1. Oncogene,
- 35



6:439-444, 1991.

25. Kallakury, B.V.S., Sheehan, C.E., Ambros, R.A.,  
Fisher, H.A.G., Kaufman, R.P., and Ross, J.S.  
5 The prognostic significance of p34<sup>cdc2</sup> and cyclic  
D1 protein expression in prostate  
adenocarcinoma. *Cancer*, 80:753-763, 1997.
26. Jiang, W., Kahn, S.M., Zhou, P., Zhang, Y.J.,  
10 Cacace, A.M., Infante, A.S., Doi, S., Santella,  
R.M., and Weinstein, I.B. Overexpression of  
cyclin D1 in rat fibroblasts causes  
abnormalities in growth control, cell cycle  
progression and gene expression. *Oncogene*,  
15 8:3447-3457, 1993.
27. Quelle, D.E., Ashmun, R.A., Shurtleff, S.A.,  
Kato, J.Y., Bar-Sagi, D., Roussel, M.F., and  
Sherr, C.J. Overexpression of mouse D-type  
20 cyclins accelerates G1 phase in rodent  
fibroblasts. *Genes Dev.*, 7:1559-1571, 1993.

#### FOURTH SERIES OF EXPERIMENTS

The INK4A gene maps to the region 9p21, and was initially described as encoding a 148 amino acid protein termed p16. The p16 protein associates exclusively with Cdk4 and Cdk6, inhibiting their complexation with D-type cyclins, and the consequent phosphorylation of pRB. This contributes to cell cycle arrest. The purpose of the present study was to evaluate patterns of p16 expression in a well characterized cohort of prostatic adenocarcinomas, while exploring potential associations between alterations of p16 and clinicopathological variables.

Normal and malignant tissues from 88 patients with prostate carcinoma were examined. In situ hybridization and immunohistochemistry assays were used to determine the status of the INK4A exon 1 $\alpha$  transcripts and levels of p16 protein, respectively. Associations between altered patterns of expression and clinicopathological variables, including pre-treatment prostate-specific antigen (PSA) level, Gleason grade, pathologic stage, and hormonal status, were evaluated using the Mantel-Haenszel chi-square test. Biochemical (PSA) relapse after surgery was evaluated using the Kaplan-Meier method and the Log rank test.

The levels of p16 expression and INK4A exon 1 $\alpha$  transcripts in normal prostate and benign hyperplastic tissues were undetectable. However, p16 nuclear overexpression was observed in 38 (43%) prostate carcinomas, while the remaining 50 (57%) cases showed undetectable p16 levels. Overexpression of p16 protein was found to correlate with increased INK4A exon 1 $\alpha$  transcripts. Moreover, p16 overexpression was associated with a higher pre-treatment PSA level ( $P=.018$ ), the use of neoadjuvant androgen ablation ( $P=.001$ ), and a sooner time to PSA relapse after

radical prostatectomy ( $P=.002$ ). These data suggest that p16 overexpression is associated with tumor recurrence and a poor clinical course in patients with prostate cancer.

5 The INK4A gene maps to the short arm of chromosome 9 (9p21), and was initially described as encoding a protein of Mr 15,845, termed p16. (1,2). The p16 protein forms binary complexes exclusively with Cdk4 and Cdk6, inhibiting their kinase activity and subsequent pRb phosphorylation  
10 during the  $G_1$  phase of the cell cycle. (1,3). Additional complexity results from the presence of a second INK4A product termed p19<sup>ARF</sup>. (4-6) The p19<sup>ARF</sup> protein has recently been shown to interact with mdm2 and to block mdm2-induced p53 degradation and transactivational silencing (7,8). The  
15 two products, p16 and p19<sup>ARF</sup>, share exons 2 and 3 of the INK4A gene, but have distinct promoters and exon 1 units, exon 1 $\alpha$  (p16) and exon 1 $\beta$  (p19<sup>ARF</sup>). The INK4A gene is mutated in a wide variety of tumor cell lines and certain primary tumors (2, 9-14). In addition, methylation of the  
20 5' CpG island of the exon 1 $\alpha$  promoter region is a frequent mechanism of p16 inactivation in primary tumors (15, 16).

In prostate cancer the role of INK4A has not been well  
25 elucidated, though analyses utilizing microsatellite markers in the vicinity of the INK4A gene have revealed loss of heterozygosity in a subset of primary and metastatic prostate tumors (17). Unlike reports of other primary tumors, INK4A inactivation, either through  
30 deletions, mutations, or through promoter methylation, appears to be an infrequent event in prostate cancer (18-23). The present study utilizes immunohistochemical and in situ hybridization assays to examine patterns of p16 expression in a well characterized cohort of prostate  
35 cancer patients treated with radical retropubic

prostatectomy. Associations between altered p16 phenotypes and clinicopathological variables were also studied to further define their potential implications in prostate cancer.

5

#### EXPERIMENTAL DETAILS

##### Experimental Details for Fourth Series of Experiments

#### 10 MATERIALS AND METHODS

Patient Characteristics and Tissues. A cohort of patients with prostatic adenocarcinoma undergoing radical prostatectomy at the Memorial Sloan-Kettering Cancer Center from 1990-1991 was retrospectively evaluated. A total of 88 patients had adequate clinical follow-up and available pathological materials. The median age at the time of surgery was 65 years (range 46 - 74 years). The median follow-up time was 64.5 months (range 10 - 94 months). Formalin-fixed, paraffin-embedded prostate tissues were obtained from the Department of Pathology. Representative hematoxylin-eosin stained sections were examined to evaluate the histopathological characteristics of each tissue section.

25

Clinicopathologic parameters examined included pre-treatment PSA, pathologic stage (24) and Gleason grade (25), both determined based on the radical prostatectomy specimen. Hormonal status of the patients was also evaluated. A portion of the cohort (34 patients - 39%) was treated with neoadjuvant androgen ablation and were defined as hormone-treated. Patients who did not receive neoadjuvant therapy were defined as hormone naive. Additionally, biochemical relapse was examined. Relapse was defined as an elevation in the serum PSA level in a

35

patient who had previously demonstrated an undetectable PSA level post-prostatectomy. That is, only patients who had an undetectable PSA level after surgery were included in the cohort, as this indicated that the surgical resection was complete and the patient was free of disease. Patients who had PSA relapse were classified as treatment failures with tumor recurrence.

Immunohistochemistry. An avidin-biotin immunoperoxidase assay was performed on formalin-fixed, paraffin-embedded tissue sections. Deparaffinized sections were treated with 1% H<sub>2</sub>O<sub>2</sub> in order to block endogenous peroxidase activity. Sections were subsequently immersed in boiling 0.01% citric acid (pH 6.0) in a microwave oven for 15 minutes to enhance antigen retrieval, allowed to cool, and incubated with 10% normal horse serum (Organon Tecknika Corp, Westchester, PA), to block non-specific tissue immunoreactivities. A well characterized antibody to p16 (Ab-1, Oncogene Research Products, Cambridge, MA; 2ug/ml final concentration) was then incubated overnight at 4°C. Biotinylated horse anti-mouse IgG antibodies (Vector Laboratories, Inc., Burlingame, CA; 1:25 final dilution) were utilized as the secondary reagents. This was followed by avidin-biotin immunoperoxidase complexes (1:25, Vector Laboratories, Inc.) for 30 minutes. Diaminobenzidine was used as the final chromogen and hematoxylin was used as the nuclear counterstain. Immunoreactivities, assessed in tissue sections from a single representative block in each case, were classified as a continuum of data from undetectable levels or (0%) to homogenous staining levels (100%). Data was independently obtained by two observers, with minor inter-observer variability which was resolved by review of the problem cases. Tumors were grouped into two categories defined as follows: Group A ( $\leq$ 5% nuclear immunoreactivity in tumor cells) and Group B ( $>$ 5% nuclear immunoreactivity

in tumor cells).

In Situ Hybridization. Primers specific for the exon 1 $\alpha$  sequence of the INK4A gene were utilized to create digoxigenin-labeled probes for in situ hybridization. Probes were cloned into a PCR-Script recombinant plasmid (Stratagene, La Jolla, CA). Plasmid DNA (1 $\mu$ g) was linearized using BamHI and XhoI. Antisense and sense riboprobes were generated from in vitro transcription of the linearized DNA using T7 and T3 RNA polymerases, respectively. Transcription was sustained for 2 hours at 37°C in 1X transcription buffer (Boehringer Mannheim, Indianapolis, IN), 20 U of RNase inhibitor, 10 mmol/L each of ATP, GTP, CTP, 6.5 mmol/L UTP and 3.5 mmol/L digoxigenin-UTP. Deparaffinized tissue sections were rinsed in water and PBS for 10 minutes. The slides were digested with Proteinase K (50  $\mu$ g/ml) for 18 minutes at 37°C in PBS, and post-fixed at 4°C in a freshly prepared solution of 4% paraformaldehyde in PBS for 5 minutes. Prehybridization was done for 30 minutes at room temperature (RT) in 50% formamide and 2X sodium chloride/sodium citrate (SSC). The hybridization buffer consisted of 50% deionized formamide (v/v), 10% dextran sulphate (50% stock solution), 2XSSC (20X stock solution), 1% SDS (10% stock solution), and 0.25 mg/ml of herring sperm DNA (10 mg/ml).

Hybridization was performed overnight at 45°C applying 10 pmol/L digoxigenin-labeled riboprobe in 50  $\mu$ l of hybridization buffer per section under a coverslip. The coverslips were removed and the slides were washed in pre-warmed 2XSSC for 20 minutes at 42°C twice, followed by washes in pre-warmed 1XSSC and 0.5XSSC at 42°C for 20 minutes. After these washes the slides were incubated in normal sheep serum diluted in buffer pH 7.5 and

successively in the same buffer with anti-digoxigenin-AP antibody (Boehringer Mannheim) at a dilution of 1:500 for 1 hour at RT. The visualization was accomplished by nitro-blue tetrazolium 5-bromo-4-chloro-3-indoylphosphate.

5 The slides were counterstained with methyl green and mounted.

10 INK4A exon-1 $\alpha$  transcript levels were examined in a subgroup of 21 cases. Consecutive tissue sections were used to analyze p16 protein by immunohistochemistry and INK4A exon-1 $\alpha$  transcript levels by in situ hybridization. As in the immunohistochemical analysis, tumors were grouped into two categories defined by the absence ( $\leq 5\%$  tumor cells with cytoplasmic staining) or presence ( $> 5\%$  tumor cells with  
15 cytoplasmic staining) of transcripts.

Statistical Methods. The statistical analyses of the data from the 88 primary prostate cancer patients were conducted as follows. The response variable, time to PSA relapse,  
20 was defined as the time from radical prostatectomy to the time of first detectable PSA measurement. Patients who did not achieve a non-measurable PSA after radical prostatectomy were excluded from the analysis. Patients who were still alive at the time of analysis without  
25 relapse were censored at the date of last follow-up. The baseline variables examined were PSA measurement at time of diagnosis, hormone status, Gleason score (hormone naïve patients only), stage of disease, and percent p16 expression.

30 Associations between p16 expression and different categorical variables (hormone status, tumor grade, tumor stage, and pretreatment PSA levels) were assessed by Mantel-Haenszel chi-square test. Continuous variables,  
35 such as pre-treatment PSA, not known to follow a particular

distribution were compared between two or more groups using Wilcoxon non-parametric tests.

5 The Cox proportional hazards model was used to examine the multivariate relationship between PSA relapse-free time from prostatectomy and the baseline variables listed above. The final model was determined using the "all subsets" procedure in SAS PHREG and the Score criterion (26). As normal and benign tissues showed little to no p16  
10 expression, positive expression was described as >5% nuclear expression. This cutpoint was specified a priori and used for the subsequent statistical analysis. Immunohistochemical and in situ hybridization studies were completed, analyzed, and recorded blind to clinical  
15 information. Kaplan-Meier estimates of relapse-free survival stratified by p16 classification were evaluated. The LIFETEST procedure in SAS was used to generate the Kaplan-Meier estimates and the resulting survival curves (26, 27). The Log rank test was used to test the  
20 hypothesis of no survival differences between p16 positive and p16 negative populations.

#### EXPERIMENTAL RESULTS

##### 25 **Experimental Results for the Fourth Series of Experiments**

The normal human prostate displayed undetectable levels of p16 protein and INK4A exon 1 $\alpha$  transcripts in ductal and acinar epithelial cells. A lack of p16 immunoreactivity in  
30 these cells was observed in hormone-treated and hormone naive cases. Fibromuscular stroma cells also showed undetectable exon 1 $\alpha$  transcripts levels. A similar negative pattern of p16 expression was observed upon the examination of prostatic tissue affected with benign  
35 hyperplasia (Figure 10).



To determine the frequency and potential clinical implications of p16 alterations in prostate cancer, we analyzed a cohort of 88 primary prostate carcinomas. Two patterns of p16 protein expression were noted. We observed that 50 of the 88 cases (57%) had very low ( $\leq 5\%$  nuclear immunoreactivity; 7 cases) or undetectable (43 cases) levels of p16 protein expression (Group A) (Figure 11A). In a subgroup of these cases we also performed in situ hybridization assays, which revealed that all cases had undetectable INK4A exon 1 $\alpha$  transcripts (Figure 11B). However, we noted that 38 of the 88 cases (43%) displayed nuclear staining with anti-p16 specific antibodies (Group B) (Figure 11C). Immunoreactivities in tumor cells were further stratified into three categories: 6% to 29% nuclear staining (n=11 cases); 30% to 59% nuclear staining (n=15 cases); and 60% to 100% nuclear staining (n=12 cases). In a subset of these patients, we also conducted in situ hybridization assays with the INK4A exon 1 $\alpha$  specific probe. All cases displaying positive immunoreactivities also displayed moderate to high levels of exon 1 $\alpha$  transcripts (Figure 11D).

Table 3 summarizes the associations between p16 phenotypes and clinicopathological variables, which were assessed by Chi-square analyses. Immunohistochemical detection of p16 was not associated with Gleason grade, described as either low (Gleason grade 4 - 6) or high (Gleason grade 7 - 10) ( $P=0.153$ ). Similarly, no association was observed between p16 nuclear expression and pathologic stage, defined as organ-confined ( $T_1$ ,  $T_2$ ) and non organ-confined ( $T_3$ ,  $T_4$ , or lymph node+) ( $P=0.087$ ). However, there was a strong association between p16 nuclear expression and pre-treatment PSA levels, based on cutoff points of  $<4$ , 4-10, and  $>10$  ng/ml ( $P=0.018$ ). A similar result was observed when PSA was assessed as a continuous variable

(P=0.01). In addition, we noted a significant correlation between p16 nuclear expression and the use of neoadjuvant androgen ablation (P=0.001).

**Table 3 Association of p16 Immunoreactivity with Tumor Grade, Hormonal Status, Tumor Stage, and Preoperative PSA Levels**  
P16 Immunoreactivity (& of patients)

|                          | Number of Subjects | ≤5%      | 5%       | p - value |
|--------------------------|--------------------|----------|----------|-----------|
| All Subjects             | 88                 | 50 (57%) | 38 (43%) |           |
| Gleason Grade*           | 82                 |          |          |           |
| <7                       | 30 (37)            | 24 (80)  | 6 (20)   | p=0.153   |
| ≥7                       | 18 (22)            | 11 (61)  | 7 (39)   |           |
| unable to evaluate       | 34 (41)            |          |          |           |
| Hormonal Status          | 88                 |          |          |           |
| hormone naive            | 54 (61)            | 40 (74)  | 14 (26)  | p=0.001   |
| hormone-treated          | 34 (39)            | 10 (29)  | 24 (71)  |           |
| Pathologic Stage         | 88                 |          |          |           |
| T <sub>2</sub>           | 53 (60)            | 34 (64)  | 19 (36)  | p=0.087   |
| ≥T <sub>3</sub>          | 35 (40)            | 16 (46)  | 19 (54)  |           |
| Pretreatment PSA (ng/ml) | 88                 |          |          |           |
| <4.0                     | 18 (20)            | 14 (78)  | 4 (22)   | p=0.018   |
| 4-10                     | 29 (33)            | 19 (66)  | 10 (34)  |           |
| >10                      | 41 (47)            | 17 (41)  | 24 (59)  |           |

\* Grading is based on the radical prostatectomy specimen. Patients who had received neoadjuvant androgen therapy were unable to be graded consistently. Six patients did not have Gleason grade information and were excluded from this analysis.

A strong association was also found between p16 nuclear overexpression and tumor recurrence, as defined by biochemical (PSA) relapse. Increasing p16 expression correlated with an increased relative hazard of relapse, suggesting a continuous relationship of the data. Overall, tumor recurrence was observed in 34 of 88 cases (39%). Thirteen of 50 cases (26%) with undetectable-to-low p16 expression (Group A) developed tumor recurrence. However, tumor recurrence was observed in 21 of 38 cases (55%) with p16 nuclear overexpression (Group B) (P=0.002) (Figure 12). Nevertheless, in a multivariate analysis adjusted for tumor grade, pre-treatment PSA, and pathologic stage,

overexpression of p16 did not contribute prognostic information over pre-treatment PSA, the strongest independent predictor of tumor recurrence.

5     Experimental Discussion for the Fourth Series of Experiments

10     Normal prostate tissues display undetectable levels of p16 protein and INK4A exon 1 $\alpha$  transcripts. It has been reported that p16 expression is low to undetectable in most normal human tissues analyzed (15, 28, 29). In support of these observations, there are relatively low and near-constant levels of p16 protein and mRNA throughout the cell cycle of normal lymphocytes in culture (30). The lack of  
15     p16 expression in hyperplastic glands parallels that of normal prostatic tissue. Based on these data, it is our hypothesis that these negative p16 phenotypes reflect basal physiologic levels of p16.

20     Primary prostatic adenocarcinomas revealed two distinct p16 phenotypes. Most tumors were found to have undetectable or very low levels of p16 protein expression (Group A - 57% of cases). This was associated with low levels or absence of INK4A exon 1 $\alpha$  transcripts. Another group of tumors showed  
25     elevated p16 protein expression (Group B - 43%) which was consistently associated with increased INK4A exon 1 $\alpha$  transcripts. These findings suggest an upregulation of the INK4A- $\alpha$  gene, resulting in p16 protein overexpression. Patients in Group B had a more aggressive course,  
30     demonstrated by high levels of pre-treatment PSA (P=.018) and a sooner time to biochemical (PSA) relapse (P=.002). A worse prognosis for Group B is also revealed by the trending association of p16 overexpression with higher pathologic stage.

35

The negative phenotype observed in Group A might correspond to the normal physiologic state, reflecting low-to-undetectable p16 levels. Alternatively, it could be related to mutations affecting the INK4A gene, especially homozygous deletions, or methylation of the INK4A exon 1 $\alpha$  promoter region. Nevertheless, it has been reported that these events are infrequent in prostate cancer (17-23). Furthermore, it appears the tumors with INK4A mutations have a more aggressive clinical course (31-33). Contrary to this, in the present study, we observed that Group A patients had a less aggressive behavior than Group B patients. For these reasons, we hypothesize that the negative phenotype observed in Group A is more likely a reflection of the normal physiologic state.

The up-regulation of the INK4A- $\alpha$  gene, resulting in the overexpression of p16 protein, may develop through different mechanisms. An association between increased p16 transcript and protein levels occur in tumor cell lines and certain primary neoplasms that lack functional pRb (1, 34-37). Moreover, p16-mediated inhibition of cell cycle progression appears to be dependent upon functional pRb (38, 39). These data support an association between p16 and pRb, where absence of functional pRb limits p16 activity and possibly promotes INK4A- $\alpha$  upregulation. Alternatively, enhanced activation of the INK4A- $\alpha$  gene may occur. E2F1, a direct activator of the INK4A exon 1 $\beta$  promoter, does not appear to directly activate INK4A- $\alpha$  transcription (40). However, evidence does exist for an indirect effect, as E2F1 overexpression has been reported to markedly increase p16 transcripts and p16-related CKI activity (41). Overexpression of cyclin D1 and/or of Cdk4 may also influence p16 expression, through a compensatory feedback loop where deregulation of cyclin D/Cdk4 complexes results in increased levels of p16 protein (28, 42). In

summary, it appears that an altered RB axis could trigger p16 overexpression in certain systems.

Cellular stress produced by replicative senescence (43-45),  
5 hyperthermia (46), and UV irradiation (47) has been  
reported to trigger p16 overexpression. In the present  
study, another type of cellular stress, androgen ablation,  
may account in part for this observed phenomenon. A subset  
of patients were treated with neoadjuvant androgen  
10 ablation, a strategy reported to decrease the incidence of  
positive surgical margins after prostatectomy (48). In the  
present study, p16 overexpression was observed in 71% of  
hormone-treated versus 26% of hormone-naive patients  
( $P=.001$ ). These data suggest that p16 expression may be  
15 enhanced by androgen depletion. Androgens are known to  
modulate the expression of other CKI's such as p27 and p21  
(49). In addition, it has been reported that the presence  
of androgens triggers downregulation of p16 in LNCaP cells  
(50), a finding consistent with our observation of p16  
20 overexpression in cases of androgen ablation. It is also  
possible that the association between p16 expression and  
androgen ablation may, in part, reflect staging bias by  
clinicians. In this setting, patients thought to have  
advanced disease may have been treated with neoadjuvant  
25 therapy.

Based on the the above referred data, it is our working  
hypothesis that p16 overexpression in prostate cancer  
represents an altered phenotype, which identifies a  
30 subgroup of patients with a higher likelihood of post  
surgical failure and tumor recurrence. In support of this  
postulate, a preliminary report in prostate cancer has  
demonstrated an association between p16 overexpression and  
poor outcome, as related to biochemical failure (51). In  
35 addition, p16 overexpression has been associated with tumor

5 progression and a poor prognosis in ovarian (52) and breast cancers (37). Though p16 acts as a negative cell cycle regulator, specific mechanisms may contribute to its altered expression, overcoming p16-mediated tumor suppressor activities. Ongoing studies may elucidate mechanisms of p16 overexpression relative to androgen depletion and/or alterations in the RB axis.

REFERENCES FOR THE FOURTH SERIES OF EXPERIMENTS.

1. Serrano M., Hannon G.J., and Beach D. "A new regulatory motif in cell cycle control causing specific inhibition of cyclinD-cdk4." Nature, 366:704-707  
5 (1993).
2. Kamb A., Gruis N., Weaver-Feldhaus J., Liu Q., Harshman K., Tavitigian S., Stockert E., Day R. 3rd, Johnson B., and Skolnick M. "A cell cycle regulator potentially involved in genesis of many tumor types." Science, 264:436-440 (1994).  
10
3. Quelle D., Ashmun R., Hannon G., Rehberger P., Trono D., Richter H., Walker C., Beach D., Sherr C., and Serrano M. "Cloning and characterization of murin p16<sup>INK4a</sup> and p15<sup>INK4b</sup> genes." Oncogene, 11:635-645 (1995).  
15
4. Quelle D., Zindy F., Ashmun R., and Sherr C. "Alternative reading frames of the INK4a tumor suppressor gene encode two unrelated proteins capable of inducing cell cycle arrest." Cell, 83:993-1000 (1995).  
20
5. Mao L., Merlo A., Bedi G., Shapiro G., Edwards C., Rollins B., and Sidransky D. "A novel p16INK4A transcript." Cancer Res., 55:2995-2997 (1995).  
25
6. Stone S., Jiang P., Dayananth P., Tavitigian S., Katcher H., Parry D., Peters G., and Kamb A. "Complex structure and regulation of the P16 (MTS1) locus." Cancer Res. 55:2988-2994 (1995).  
30
7. Pomerantz J., Schreiber-Agus N., Liegeois N.J., Silverman A., Alland L., Chin L., Potes J., Chen K., Orlow I., Lee H.W., Cordon-Cardo C., and DePinho R.  
35

"The Ink4a tumor suppressor gene product, p19<sup>Arf</sup>, interacts with MDM2 and neutralizes MDM2's inhibition of p53." Cell, 92:713-723 (1998).

- 5      8.    Zhang Y., Xiong Y., and Yarbrough W. "ARF promotes MDM2 degradation and stabilized p53: ARF-INK4a locus deletion impairs both the Rb and p53 tumor suppression pathways." Cell, 92:725-734 (1998).
- 10     9.    Jen J., Harper J., Bigner S., Bigner D., Papadopoulos N., Markowitz S., Willson J., Kinzler K., and Vogelstein B. "Deletion of p16 and p15 genes in brain tumors." Cancer Res. 54:6353-6358 (1994).
- 15     10.   Spruck C. III, Gonzalez-Zulueta M., Shibata A., Simoneau A., Lin M., Gonzales F., Tsai Y., and Jones P. "p16 gene in uncultured tumours." Nature, 370:183-184 (1994).
- 20     11.   Orlow I., Lacombe L., Hannon G., Serrano M., Pellicer I., Dalbagni G., Reuter V., Zhang Z., Beach D., and Cordon-Cardo C. "Deletion of the p16 and p15 genes in human bladder tumors." J. Natl. Cancer Inst., 87:1524-1529 (1995).
- 25     12.   Reed A., Califano J., Cairns P., Westra W., Jones R., Koch W., Ahrendt S., Eby Y., Sewell D., Nawroz H., Bartek J., and Sidransky D. "High frequency of p16 (CDKN2/MTS1/INK4A) inactivation in head and neck squamous cell carcinoma." Cancer Res., 56:3630-3633 (1996).
- 30     13.   Washimi O., Nagatake M., Osada H., Ueda R., Koshikawa T., Seki T., Takahashi T., and Takahashi T. "In vivo occurrence of p16 (MTS1) and p15 (MTS2) alterations
- 35



preferentially in non-small cell lung cancers" Cancer Res., 55:514-517 (1995).

14. Takeuchi S., Bartram C., Seriu T., Miller C., Tobler  
5 A., Janssen J., Reiter A., Ludwig W., Zimmermann M.,  
Schwaller J., Lee E., Miyoshi I., and Koeffler HP.  
"Analysis of a family of cyclin-dependent kinase  
inhibitors: p15/MTS2/INK4B, p16/MTS1/INK4A, and p18  
10 Blood, 86:755-760 (1995).
15. Gonzales-Zulueta M, Bender C, Yang A, Nguyen T, Beart  
R, Van Tornout J, and Jones P. "Methylation of the 5'  
CpG island of the p16/CDKN2 tumor suppressor gene in  
15 normal and transformed human tissues correlates with  
gene silencing." Cancer Res., 55:4531-4535 (1995).
16. Merlo A, Herman J, Mao L, Lee D, Gabrielson E, Burger  
P, Baylin S, and Sidransky D. "5' CpG island  
20 methylation is associated with transcriptional  
silencing of the tumor suppressor p16/CDKN2/Mts1 in  
human cancers." Nature Med., 1:686-692 (1995).
17. Jarrard D, Bova S, Ewing C, Pin S, Nguyen S, Baylin S,  
25 Cairns P, Sidransky D, Herman J, and Isaacs W.  
"Deletional, mutational, and methylation analyses of  
CDKN2 (p16/MTS1) in primary and metastatic prostate  
cancer." Genes, Chrom. & Cancer, 19:90-96 (1997).
18. Mangold K, Takahashi H, Brandigi C, Wada T, Wakui S,  
30 Furusato M, Boyd J, Chandler F, and Allsbrook W. "p16  
(CDKN2/MTS1) gene deletions are rare in prostatic  
carcinomas in the United States and Japan." J. Urol.,  
157:1117-1120 (1997).

19. Park D, Wilczynski S, Pham E, Miller C, and Koeffler HP. "Molecular analysis of the INK4 family of genes in prostate carcinomas." J. Urol., 157: 1995-1999, (1997).
20. Tamimi Y, Bringuier P, Smit F, van Bokhoven A, Debruyne F, and Schalken J. "p16 mutations/deletions are not frequent events in prostate cancer." Brit. J. Cancer, 74:120-122 (1996).
21. Chen W, Weghorst M, Sabourin C, Wang Y, Wang D, Bostwick D, and Stoner G. "Absence of p16/MTS1 gene mutations in human prostate cancer." Carcinogenesis, 17: 2603-7 (1996).
22. Jarrard D, Bova S, Ewing C, Pin S, Nguyen S, Baylin S, Cairns P, Sidransky D, Herman J, and Isaacs W. "Deletional, mutational, and methylation analyses of CDKN2 (p16/MTS1) in primary and metastatic prostate cancer." Genes, Chrom. & Cancer, 19:90-96 (1997).
23. Gu K, Mes-Masson AM, Gauthier J, and Saad F. "Analysis of the p16 tumor suppressor gene in early-stage prostate cancer." Mol. Carcinogenesis, 21:164-170 (1998).
24. Schroder F, Hermanek P, Denis L, Fair W, Gospodarowicz M, and Pavone-Macaluso M. "The TNM classification of prostate cancer." Prostate, 4:129-38 (1992).
25. Gleason D and Mellinger G. "Prediction of prognosis for prostatic adenocarcinoma by combined histological grading and clinical staging." J. Urol, 111:58-64 (1974).

26. SAS Institute Inc. SAS/STAT User Guide, Version 6.  
Cary, NC: SAS Institute Inc. (1990).
- 5 27. Kaplan EL, Meier P. Nonparametric estimation from  
incomplete observations. *JASA*, 53:457-481 (1958).
- 10 28. Yao J, Pollock R, Lang A, Tan M, Pisters W, Goodrich  
D, El-Naggar A, and Yu D. "Infrequent mutation of the  
p16/MTS1 gene and overexpression of cyclin-dependent  
kinase 4 in human primary soft-tissue sarcoma." Clin.  
Cancer Res., 4:1065-1070 (1998).
- 15 29. Shapiro G, Edwards C, Kobzik L, Lodleski J, Richards  
W, Sugarbaker D, and Rollins B. "Reciprocal Rb  
inactivation and p16INK4 expression in primary lung  
cancers and cell lines." Cancer Res., 55:505-509  
(1995).
- 20 30. Soucek T, Pusch O, Hengstschlager-Ottnd E, Wawra E,  
Bernaschek G, and Hengstschlager M. "Expression of  
the cyclin-dependent kinase inhibitor p16 during the  
ongoing cell cycle." FEBS Letters, 373:164-169  
(1995).
- 25 31. Taga S, Osaki T, Ohgami A, Imoto H, Yoshimatsu T,  
Yoshino I, Yano K, Nakanishi R, Ichiyoshi Y, and  
Yasumoto K. "Prognostic value of the  
immunohistochemical detection of p16INK4 expression in  
nonsmall cell lung carcinoma." Cancer, 80:389-95  
30 (1997).
- 35 32. Garcia-Sanz R, Gonzalez M, Vargas M, Chillon M,  
Balanzategui A, Barbon M, Flores M, and San Miguel J.  
"Deletions and rearrangements of cyclin-dependent  
kinase 4 inhibitor gene p16 are associated with poor

prognosis in B cell non-Hodgkin's lymphomas." Leukemia, 11: 1915-20, (1997).

- 5           33. Straume O and Akslen L. "Alterations and prognostic significance of p16 and p53 protein expression in subgroups of cutaneous melanoma." Int. J. Cancer 74:535-539 (1997).
- 10           34. Parry D, Bates S, Mann DJ, and Peters G. "Lack of cyclin D-cdk complexes in Rb-negative cells correlates with high levels of p16INK4/MTS1 tumour suppressor gene product." EMBO J., 14:503-511 (1995).
- 15           35. Ueki K, Ono Y, Henson JW, Efird JT, von Deimling A, Louis DN. "CDKN2/p16 or RB alterations occur in the majority of glioblastomas and are inversely correlated." Cancer Res., 56:150-153 (1996).
- 20           36. Li Y, Nichols M, Shay J, and Xiong Y. "Transcriptional repression of the D-type cyclin-dependent kinase inhibitor p16 by the retinoblastoma susceptibility gene product pRb." Cancer Res., 54:6078-6082 (1994).
- 25           37. Dublin E, Patel N, Gillett C, Smith P, Peters G, and Barnes D. "Retinoblastoma and p16 proteins in mammary carcinoma: their relationship to cyclin D1 and histopathological parameters." Int. J. Cancer, 79:71-5 (1998).
- 30           38. Lukas J, Parry D, Aagaard L, Mann DJ, Bartkova J, Strauss M, Peters G, and Bartek J. "Retinoblastoma-protein-dependent-cell-cycle inhibition by the tumour suppressor p16." Nature, 375:503-506 (1995).

39. Craig C, Kim M, Ohri E, Wersto R, Katayose D, Li Z, Choi Y, Mudahar B, Srivastava S, Seth P, and Cowan K. "Effects of adenovirus-mediated p16INK4A expression on cell cycle arrest are determined by endogenous p16 and Rb status in human cancer cells." Oncogene, 16:265-272 (1998).
40. Robertson K and Jones P. "The human ARF cell cycle regulatory gene promoter is a CpG island which can be silenced by DNA methylation and down-regulated by wild-type p53." Mol. Cell. Biol., 18:6457-6473 (1998).
41. Khleif S, DeGregori J, Yee C, Otterson G, Kaye F, Nevins J, and Howley P. "Inhibition of cyclin D-CDK4/CDK6 activity is associated with an E2F-mediated induction of cyclin kinase inhibitor activity." Proc. Natl. Acad. Sci. USA, 93:4350-4354 (1996).
42. Burns K, Ueki K, Jhung S, Koh J, and Louis D. "Molecular genetic correlates of p16 cdk4, and pRb immunohistochemistry in glioblastomas." J. Neuropathol. & Exp. Neurology, 57:122-130 (1998).
43. Hara E, Smith R, Parry D, Tahara H. Stone S, and Peters G. "Regulation of p16 CDKN2 expression and its implications for cell immortalization and senescence." Mol. Cell. Biol., 16:859-867 (1996).
44. Palmero I, McConnel B, Parry D, Brookes S, Hara E, Bates S, Jat P, and Peters G. "Accumulation of p16INK4a in mouse fibroblasts as a function of replicative senescence and not of retinoblastoma gene status." Oncogene, 15:495-503 (1997).

45. Yeager T, DeVries S, Jarrard D, Kao C, Nakada S, Moon T, Bruskewitz R, Stadler W, Meisner L, Gilchrist K, Newton M, Waldman F, and Reznikoff C. "Overcoming cellular senescence in human cancer pathogenesis." Genes & Devel., 12:163-174 (1998).  
5
46. Valenzuela MT, Nunez MI, Villalobos M, Siles E, McMillan T, Pedraza V, and Almodovar R. "A comparison of p53 and p16 expression in human tumor cells treated with hyperthermia or ionizing radiation." Int. J. Cancer, 72:307-312 (1997).  
10
47. Wang X, Gabrielli B, Milligan A, Dickinson J, Antalis T, and Ellem K. "Accumulation of p16<sup>CDKN2A</sup> in response to ultraviolet irradiation correlates with a late S-G<sub>2</sub>-phase cell cycle delay." Cancer Res., 56:2510-2514 (1996).  
15
48. Witjes W, Schulman C, and Debruyne F. "Preliminary results of a prospective randomized study comparing radical prostatectomy versus radical prostatectomy associated with neoadjuvant hormonal combination therapy in T2-3 N0 M0 prostatic carcinoma. The European Study Group on Neoadjuvant Treatment of Prostate Cancer." Urology, 49S:65-9 (1997).  
20  
25
49. Kokontis J, Hay N, and Liao S. "Progression of LNCaP prostate tumor cells during androgen deprivation: Hormone-independent growth, repression of proliferation by androgen, and role for p27<sup>Kip1</sup> in androgen-induced cell cycle arrest." Mol. Endocrinol., 12:941-953 (1998).  
30
50. Lu S, Tsai SY, and Tsai M-J. "Regulation of androgen-dependent prostatic cancer cell growth: androgen  
35

regulation of CDK2, CDK4, and CKI p16 genes." Cancer Res., 57:4511-4516 (1997).

51. Halvorsen OJ, Hostmark J, Haukass, Hoisaeter P, and  
5 Akslen L. "Prognostic importance of p16 and CDK4  
proteins in localized prostate cancer." Proc. Am.  
Assoc. Cancer Res., 38:526 (1997).
52. Dong Y, Walsh M, McGuckin M, Gabrielli B, Cummings M,  
10 Wright RG, Hurst T, Khoo SK, and Parsons P.  
"Increased expression of cyclin-dependent kinase  
inhibitor 2 (CDKN2A) gene product p16<sup>ink4A</sup> in ovarian  
cancer is associated with progression and unfavorable  
prognosis." Int. J. Cancer, 74:57-63 (1997).

15

**What is Claimed is:**

1. A method for determining the aggressiveness of a prostate carcinoma comprising:
  - 5 (a) obtaining a sample of the prostate carcinoma; and
  - (b) detecting the presence of p27 protein in the prostate carcinoma, the absence of p27 indicating that the prostate carcinoma is aggressive.
- 10 2. A method for diagnosing a benign prostate hyperplasia comprising:
  - 15 (a) obtaining an appropriate sample of the hyperplasia; and
  - (b) detecting the presence of the p27 RNA, a decrease of the p27 RNA indicating that the hyperplasia is benign.
- 20 3. A method of claim 2, further comprising detecting the protein expression of p27 wherein this additional step may be performed before or after the detection of the presence of the p27 RNA.
- 25 4. A method for predicting the life-span of patient with prostate carcinoma comprising:
  - (a) obtaining a sample of the prostate carcinoma; and
  - 30 (b) detecting the presence of p27 protein in the prostate carcinoma, the presence of the p27 protein indicating that the patient can live longer than the patient who are undetectable p27 protein.
- 35 5. A method for increasing the life-span of patient with



prostate carcinoma comprising inducing the expression of p27 protein in the prostate carcinoma.

- 5 6. A method for prolong life-span of patient with prostate carcinoma which comprises introducing a nucleic acid molecule having sequence encoding a p27 protein into the carcinoma cell under conditions permitting expression of said gene so as to prolong the life-span of the patient with said prostate carcinoma.  
10
7. The method of claim 6, wherein the nucleic acid molecule comprises a vector.
- 15 8. The method of claim 7, wherein the vector is an adenovirus vector, adenoassociated virus vector, Epstein-Barr virus vector, retrovirus vector or vaccinia virus vector.
- 20 9. A method for prolong life-span of patient with prostate carcinoma which comprises introducing an effective amount of p27 protein into the carcinoma cell so as to thereby prolong the life-span of the patient with said prostate carcinoma.  
25
10. A method for prolong life-span of patient with prostate carcinoma which comprises introducing an effective amount of a substance capable of stabilizing the p27 protein into the carcinoma cell so as to  
30 thereby prolong the life-span of the patient with said prostate carcinoma.
11. A composition for prolong life-span of patient with prostate carcinoma which comprises an effective amount  
35 of a nucleic acid molecule having sequence encoding a

p27 protein and a suitable carrier.

12. A composition for prolong life-span of patient with prostate carcinoma which comprises an effective amount of the p27 protein and a suitable carrier.
13. A composition for prolong life-span of patient with prostate carcinoma which comprises an effective amount a substance capable of stabilizing the p27 protein and a suitable carrier.
14. A method for determining the rate of proliferation of a prostate cancer comprising:
  - (a) obtaining a sample of the prostate cancer; and
  - (b) detecting the presence of p21 protein in the prostate cancer, the presence of p21 indicating that the prostate cancer will have a high proliferation rate.
15. A method for determining the rate of proliferation of a prostate cancer comprising:
  - (a) obtaining a sample of the prostate cancer; and
  - (b) detecting the mdm2 expression in the prostate cancer, the overexpression of mdm2 indicating that the prostate cancer will have high proliferation rate.
16. A method for determining whether a prostate cancer would be metastatic comprising:
  - (a) obtaining a sample of the prostate cancer; and
  - (b) detecting the level of cyclin D1 expression in the prostate cancer, the overexpression of cyclin D1 indicating that the prostate cancer will be metastatic.

17. The method of claim 16, wherein the prostate cancer is metastatic to bone.

18. A method for determining the tumor recurrence in prostate cancer comprising:

5

- (a) obtaining a sample of the prostate cancer; and
- (b) detecting the expression of the cyclin-dependent kinase inhibitor p16 in the prostate cancer, the overexpression of p16 indicating that the prostate cancer will have high tumor recurrence.

10

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199

MARKERS FOR PROSTATE CANCER

Abstract of the Disclosure

5     This invention provides a method for determining the  
aggressiveness of a prostate carcinoma comprising: (a)  
obtaining a sample of the prostate carcinoma; and (b)  
detecting the presence of p27 protein in the prostate  
carcinoma, the absence of p27 indicating that the prostate  
10    carcinoma is aggressive. This invention also provides a  
method for diagnosing a beign prostate hyperplasia  
comprising: (a) obtaining an appropriate sample of the  
hyperplasia; and (b) detecting the presence of the p27 RNA,  
a decrease of the p27 RNA indicating that the hyperplasia  
15    is beign. This invention provides various uses of p27 in  
prostate cancer. Finally, this invention also provides  
different marker for prostate cancer.




FIG. 1H



FIG. 1G



FIG. 2A




FIG. 2B




FIG. 2C









FIG. 2D



0.0015  $E/K^2$    
0.003  $E/K^2$    
0.006  $E/K^2$    
0.012  $E/K^2$  






FIG. 4A



FIG. 4B



FIG. 4C



FIG. 4D



FIG. 4E




FIG. 4F

FIG. 5A




FIG. 5B





FIG. 5C



FIG. 5D




**FIG. 7A****FIG. 7B****FIG. 7C**

FIG. 8A




FIG. 8B




FIG. 9C



FIG. 9B




FIG. 9A



FIG. 9F



FIG. 9E



FIG. 9D



FIG. 10B



FIG. 10A




FIG. 11B




FIG. 11D




FIG. 11A




FIG. 11C

